arXiv:math/0308178v1 [math.AG] 19 Aug 2003

FULL TWISTS FACTORIZATION FORMULA FOR DOUBLE NUMBER OF STRINGS

VIK. S. KULIKOV

Abstract. A formula for factorizations of the full twist in the braid group Br_{2m} depending on any four factorizations of the full twist in Br_m is given. Applying this formula, a symplectic 4-manifold X and two isotopic generic coverings $f_i : X \to \mathbb{CP}^2$, branched, respectively, along cuspidal Hurwitz curves $\bar{H}_i \subset \mathbb{CP}^2$ (without negative nodes) having different braid monodromy factorization types, are constructed. The class of the fundamental groups of the complements of the affine plane Hurwitz curves is described in terms of generators and defining relations.

0. Introduction

Let (X, L) be a polarized projective surface over the field $\mathbb{C}$, where the polarization L is a very ample line bundle on X. Three generic sections of L define a generic covering $f : X \to \mathbb{CP}^2$ branched over an algebraic cuspidal curve $\bar{H} \subset \mathbb{CP}^2$ (possibly, one should take three sections of $L^{\otimes 2}$ instead of the sections of L to obtain a generic covering of the plane ([Ku-Ku])). Recently, Auroux and Katzarkov (see [Au], [Au-Ka]) obtained a similar result in symplectic case. More precisely, let (X, ω) be a compact symplectic 4-manifold with symplectic form ω whose class $[\omega] \in H^2(X, \mathbb{Z})$. Fix an ω-compatible almost complex structure J and corresponding Riemannian metric g. Let L be a line bundle on X whose first Chern class is $[\omega]$. Then, for $k >> 0$, the line bundle $L^{\otimes k}$ admits many approximately holomorphic sections so that one can choose three of them which give an approximately holomorphic generic covering $f_k : X \to \mathbb{CP}^2$ of degree $N_k = k^2\omega^2$ branched over a cuspidal Hurwitz curve $\bar{H}_k$ (the notion of Hurwitz curves is a generalization of the notion of plane algebraic curves; see the definition in section 5). In algebraic case, if $\deg f \geq 12$, then f is determined uniquely by $\bar{H}$ ([Ku], [Nem]). Any generic covering $f : X \to \mathbb{CP}^2$ of degree N branched over a cuspidal Hurwitz curve $\bar{H}$ determines

---

Date: This version: July 2003.

The work was partially supported by RFBR (No. 02-01-00786) and INTAS (No. 00-0269).

(and is determined by) its monodromy $\overline{\mu}$, that is, an epimorphism $\overline{\mu} : \pi_1(\mathbb{CP}^2 \setminus \bar{H}) \to \mathfrak{S}_N$ to the symmetric group $\mathfrak{S}_N$. Therefore one of the main problems is to investigate the properties of the fundamental groups of the complements of Hurwitz curves in order to have a possibility to construct interesting examples of symplectic 4-manifolds.

In [Ku1], a class $\mathcal{C}$ of groups, called C-groups, was defined. This class coincides (see Proposition 1.3) with a class consisting of the groups which are given by finite presentations of the following form: for some integer m, a function $h : \{1, \dots, m\} \to \mathbb{Z}$, and a subset $W = \{w_{i,j,k} \in \mathbb{F}_m \mid 1 \leq i, j \leq m,\ 1 \leq k \leq h(i, j)\} \subset \mathbb{F}_m$, where $\mathbb{F}_m$ is the free group generated by an alphabet $\{x_1, \dots, x_m\}$, a group $G \in \mathcal{C}$ possesses the presentation

$$G_W =< x_1, \dots, x_m \mid x_i = w_{i,j,k}^{-1} x_j w_{i,j,k},\ w_{i,j,k} \in W > . \qquad (1)$$

Let $\varphi_W : \mathbb{F}_m \to G_W$ be the canonical epimorphism. The elements $\varphi_W(x_i) \in G$, $1 \leq g \leq m$, and the elements conjugated to them are called the *C-generators* of the C-group G. Let $f : G_1 \to G_2$ be a homomorphism of C-groups. It is called a *C-homomorphism* if the images of the C-generators of G_1 under f are C-generators of the C-group G_2. We will distinguish the groups $G \in \mathcal{C}$ up to C-isomorphisms. Note that the class $\mathcal{C}$ contains the subclasses $\mathcal{K}$ and $\mathcal{L}$, respectively, of the knot and link groups given by Wirtinger presentation. In [Ku1], it was proved that the class $\mathcal{C}$ coincides with the class of the fundamental groups of the complements of orientable closed surfaces in the 4-dimensional sphere S^4 (with generalized Wirtinger presentation).

We prove (Theorem 6.1) that the class $\mathcal{H} = \{\, \pi_1(\mathbb{C}^2 \setminus H) \,\}$ of the fundamental groups of the complements of affine Hurwitz curves H (that is, $H = \bar{H} \cap (\mathbb{CP}^2 \setminus L_\infty)$, where $\bar{H}$ is a Hurwitz curve in $\mathbb{CP}^2$ and L_∞ is a projective line in $\mathbb{CP}^2$ in general position with respect to $\bar{H}$) coincides with a subclass of $\mathcal{C}$ consisting of the C-groups G_W with presentation (1) such that the set of the set of words $\{[x_i, x_1 \dots x_m]\}_{i=1,\dots,m}$ is a subset of W. We also describe (Theorem 6.3) the intersection $\mathcal{H} \cap \mathcal{K}$ of the classes $\mathcal{H}$ and $\mathcal{K}$. Besides, we prove (Corollary 1.15) that the class $\mathcal{C}$ coincides with a class $\mathcal{C}_{Br}$ of groups which possess presentations of the form

$$G_B =< x_1, \dots, x_m \mid x_i^{-1} b(x_i) = \mathbf{1},\ i = 1, \dots, m,\ b \in B >$$

for some $m \in \mathbb{N}$ and a finitely generated subgroup B of the braid group Br_m, where $b(x_i)$ is the image of x_i under the standard action of $b \in Br_m$ on $\mathbb{F}_m$. The proof of these results is based on a full twist factorizations formula in the braid group Br_{2m} (see section 3). This formula also is applied to prove the existence of a symplectic 4-manifold

X and two isotopic generic coverings $f_i : X \to \mathbb{CP}^2$, $i = 1, 2$, branched, respectively, over cuspidal Hurwitz curves (without negative nodes) $\bar{H}_i \subset \mathbb{CP}^2$ having different braid monodromy factorization types (see Theorem 7.7).

1. C-GROUPS

Let $\mathbb{F}$ be the free group generated by an alphabet $\{x_j \mid j \in \mathbb{N}\}$. Below, denote by $\mathbb{F}_m$ the subgroup of $\mathbb{F}$ generated by $x_1, \dots, x_m$, and $\mathbb{F}_{m,k}$ the subgroup of $\mathbb{F}$ generated by $x_{k+1}, \dots, x_{k+m}$. Notation $w(x_1, \dots, x_m)$ means a word in letters $x_1, \dots, x_m$ and their inverses considered as an element in the group $\mathbb{F}_m$.

Let W be a subset of $\mathbb{F}_m$ and $N_m(W)$ the normal closure of W in $\mathbb{F}_m$. Denote by $\varphi_W : \mathbb{F}_m \to \mathbb{F}_m / N_m(W) = G(W)$ the canonical epimorphism. The set

$$R_W = \{w_j(x_1, \dots, x_m) = \mathbf{1} \mid w_j \in W\}$$

is called a *set of relations defining* $G(W)$. We say that the relations R_W imply a relation $\{w(x_1, \dots, x_m) = \mathbf{1}\}$ if $w \in N_m(W)$. Sometimes the set R_W will be written in the form

$$R_W = \{u_j(x_1, \dots, x_m) = v_j(x_1, \dots, x_m) \mid w_j(\bar{x}) = u_j(\bar{x}) v_j(\bar{x})^{-1} \in W\}$$

for some presentation of the words $w_j \in W$ in the form $w_j = u_j v_j^{-1}$.

Let $W_1 \subset \mathbb{F}_{m_1}$, $W_2 \subset \mathbb{F}_{m_2}$, and $f : G(W_1) \to G(W_2)$ be a homomorphism. We say that f is the *canonical homomorphism* if

$$f(\varphi_{W_1}(x_i)) = \varphi_{W_2}(x_i)$$

for all $i \leq \min(m_1, m_2)$. In particular, if $W_1 \subset W_2$ are two subsets in $\mathbb{F}_m$, then there is the canonical epimorphism

$$\psi_{W_1,W_2} : G(W_1) \to G(W_2) = G(W_1)/N_{G(W_1)}(\varphi_{W_1}(W_2 \setminus W_1)), \qquad (2)$$

where $N_{G(W_1)}(\varphi_{W_1}(W_2 \setminus W_1))$ is the normal closure in $G(W_1)$ of the set $\varphi_{W_1}(W_2 \setminus W_1)$.

We say that two sets of relations

$$R_{W_i} = \{w_{i,j} = \mathbf{1} \mid w_{i,j} \in W_i\}, \qquad i = 1, 2,$$

are *equivalent* if $N_m(W_1) = N_m(W_2)$. More generally, for $k \leq m$, let $W_1 \subset \mathbb{F}_k$ and $W_2 \subset \mathbb{F}_m$ be two subsets of elements in $\mathbb{F}_k$ and $\mathbb{F}_m$, respectively. The sets of relations

$$R_{W_1} = \{w_{1,j} = \mathbf{1} \mid w_{1,j} \in W_1\}$$

and

$$R_{W_2} = \{w_{2,j} = \mathbf{1} \mid w_{2,j} \in W_2\}$$

are called *equivalent* if there is the canonical isomorphism $h : G(W_1) \to G(W_2)$.

Claim 1.1. *For $W_1, W_2 \subset \mathbb{F}_k$ and $\overline{W}_1 \subset \mathbb{F}_m$, $m \geq k$, if the sets of relations R_{W_1} and $R_{\overline{W}_1}$ are equivalent, then the sets of relations $R_{W_1 \cup W_2}$ and $R_{\overline{W}_1 \cup W_2}$ are equivalent.*

Proof. Straightforward. □

Claim 1.2. *Let $w(x_1, \dots, x_m) = x_{j_1}^{\varepsilon_1} \dots x_{j_n}^{\varepsilon_n}$, $\varepsilon_k = \pm 1$ for $k = 1, \dots, n$, be a word in the letters $x_1, \dots, x_m$ and their inverses of the letter length $n \geq 2$. Then for any pair $i_1, i_2 \in \{1, \dots, m\}$, the relation $\{x_{i_1} = w^{-1} x_{i_2} w\}$ in $\mathbb{F}_m$ is equivalent to the union of the sets of relations*

$$\begin{array}{rl} R_w = & \{x_{m+k} = x_{j_k}^{-\varepsilon_k} x_{m+k-1} x_{j_k}^{\varepsilon_k} \mid k = 2, \dots, n-1\} \cup \\ & \{x_{m+1} = x_{j_1}^{-\varepsilon_1} x_{i_2} x_{j_1}^{\varepsilon_1},\ x_{i_1} = x_{j_n}^{-\varepsilon_n} x_{m+n-1} x_{j_n}^{\varepsilon_n}\} \end{array}$$

in $\mathbb{F}_{m+n-1}$.

Proof. Straightforward. □

Let I be any subset in $\{1, \dots, N\}^3$. Since for $(i, j, k) \in \{1, \dots, N\}^3$ the relations $x_i = x_j^{-1} x_k x_j$ and $x_k = x_j x_i x_j^{-1}$ are equivalent, Claims 1.1 and 1.2 imply the following proposition.

Proposition 1.3. *Any C-group G is canonically C-isomorphic to a C-group $\overline{G}$ with C-presentation of the form*

$$\overline{G} \simeq < x_1, \dots, x_N \mid x_i = x_j^{-1} x_k x_j \text{ for } (i, j, k) \in I >,$$

i.e., the definition of C-groups given in Introduction coincides with the definition of C-groups given in [Ku1].

Note that the relation $x_i = x_j$ is a C-relation, since it can be written in the form $x_i = x_j^{-1} x_j x_j$.

Example. Let Br_m be the braid group on m strings. It is generated by generators $a_1, \dots, a_{m-1}$ being subject to the relations

$$\begin{array}{lll} a_i a_{i+1} a_i & = a_{i+1} a_i a_{i+1} & 1 \leq i \leq m-1, \\ a_i a_k & = a_k a_i & \mid i - k \mid \geq 2. \end{array} \tag{3}$$

The braid group Br_m with presentation (3) possesses a natural structure of C-group. Indeed, let

$$\begin{array}{l} R_{W(Br_m)} = \{ x_{2i+1} = x_{2i-1}^{-1} x_{2i} x_{2i-1} \mid i = 1, \dots, m-1 \} \cup \\ \{ x_{2i-1} = x_{2i}^{-1} x_{2i+1} x_{2i} \mid i = 1, \dots, m-1 \} \cup \\ \{ x_{2i-1} = x_{2j-1}^{-1} x_{2i-1} x_{2j-1} \mid |i-j| \geq 2,\ 1 \leq i, j \leq m-1 \} \end{array}$$

be a set of C-relations in $\mathbb{F}_{2m-3}$. Put $G_{Br_m} = G(W(Br_m))$. One can check that the homomorphism $f : Br_m \to G_{Br_m}$ given by $f(a_i) = \varphi_W(x_{2i-1})$, $i = 1, \dots, m-1$, is an isomorphism.

For a set $W = W_f = \{w_j(x_1, \dots, x_m) \in \mathbb{F}_m \mid j \in J\}$, where $f : J \to \mathbb{F}_m$ is a map from a set J and $f(j)$ is denoted by w_j for $j \in J$, the set

$$\mathrm{sh}\,(W) = \mathrm{sh}_m\,(W) = \{w_j(x_{m+1}, \dots, x_{2m}) \mid j \in J\}$$

is called the *shift* of W by m. Denote by

$$W_{mod\,m} = \{x_i^{-1}x_{m+i} \mid i = 1, \dots, m\}$$

and call the sets

$$d(W) = d_m(W) = W \cup \mathrm{sh}\,(W) \cup W_{mod\,m} \tag{4}$$

and

$$\overline{d}(W) = \overline{d}_m(W) = \mathrm{sh}\,(W) \cup W_{mod\,m} \tag{5}$$

in $\mathbb{F}_{2m}$, respectively, the *doubling* and the *reduced doubling of the set* W.

The doubling (respectively, the reduced doubling) of a set W can be iterated if we put

$$d^n(W) = d_{2^{n-1}m}(d^{n-1}(W)) \in \mathbb{F}_{2^n m}. \tag{6}$$

Claim 1.4. *For any $n \in \mathbb{N}$ the sets of relations R_W, $R_{d^n(W)}$, and $R_{\overline{d}^n(W)}$ are equivalent.*

Proof. Straightforward. ☐

Let B be a subgroup of $\mathrm{Aut}\,(\mathbb{F}_m)$. Denote by $R_{W(B,\mathbb{F}_m)}$ the set of relations

$$R_{W(B,\mathbb{F}_m)} = \{wg(w)^{-1} = \mathbf{1} \mid w \in \mathbb{F}_m,\ g \in B\}.$$

Claim 1.5. *Let $y_1, \dots, y_m \in \mathbb{F}_m$ be elements generating $\mathbb{F}_m$. For a subgroup $B \subset Aut\,(\mathbb{F}_m)$, the set of relations $R_{W(B,\mathbb{F}_m)}$ is equivalent to*

$$R_{W(B,\overline{y})} = \{y_i g(y_i)^{-1} = \mathbf{1} \mid 1 \le i \le m,\ g \in B\}.$$

Proof. Note that to prove Claim 1.5, it is sufficient to show that the condition that elements $w_1g(w_1)^{-1}$ and $w_2g(w_2)^{-1}$ belong to $N_m(W(B,\overline{y}))$ for some $w_1, w_2 \in \mathbb{F}_m$ implies

$$w_1^{-1}g(w_1^{-1})^{-1} \in N_m(W(B,\overline{y}))$$

and

$$w_1w_2g(w_1w_2)^{-1} \in N_m(W(B,\overline{y})).$$

If $w_1g(w_1)^{-1} \in N_m(W(B,\overline{y}))$, then

$$(w_1g(w_1)^{-1})^{-1} = g(w_1)w_1^{-1} \in N_m(W(B,\overline{y}))$$

and therefore

$$w_1^{-1}g(w_1^{-1})^{-1} = w_1^{-1}(g(w_1)w_1^{-1})w_1 \in N_m(W(B,\overline{y})).$$

If $w_1g(w_1)^{-1}, w_2g(w_2)^{-1} \in N_m(W(B,\overline{y}))$, then

$$\begin{array}{l} w_1w_2g(w_1w_2)^{-1} = w_1w_2g(w_2)^{-1}g(w_1)^{-1} = \\ w_1(w_2g(w_2)^{-1})w_1^{-1}(w_1g(w_1)^{-1}) \in N_m(W(B,\overline{y})). \end{array}$$

□

Claim 1.6. *If $B \subset Aut(\mathbb{F}_m)$ is generated by elements $b_1,\dots,b_n$ and elements $y_1,\dots,y_m$ generate $\mathbb{F}_m$, then $R_{W(B,\mathbb{F}_m)}$ is equivalent to*

$$R_{W(\overline{b},\overline{y})} = \{y_ib_j(y_i)^{-1} = \mathbf{1} \ \mid \ 1 \leq i \leq m,\ 1 \leq j \leq n\}.$$

Proof. It follows from the proof of Claim 1.5 that to prove Claim 1.6, it is sufficient to show that the condition that the elements $wg_1(w)^{-1}$ and $wg_2(w)^{-1}$ belong to $N_m(W(\overline{b},\overline{y}))$ for some $g_1,g_2 \in B$ and any $w \in \mathbb{F}_m$ implies $w(g_1^{-1}(w))^{-1} \in N_m(W(\overline{b},\overline{y}))$ and $w(g_1g_2(w))^{-1} \in N_m(W(\overline{b},\overline{y}))$.

If $wg_1(w)^{-1} \in N_m(W(\overline{b},\overline{y}))$ for all $w \in \mathbb{F}_m$, then

$$\begin{array}{l} (w(g_1^{-1}(w))^{-1})^{-1} = (g_1^{-1}(w))w^{-1} = \\ (g_1^{-1}(w))(g_1(g_1^{-1}(w)))^{-1} \in N_m(W(\overline{b},\overline{y})). \end{array}$$

If $w(g_1(w))^{-1}, w(g_2(w))^{-1} \in N_m(W(\overline{b},\overline{y}))$ for all $w \in \mathbb{F}_m$, then

$$\begin{array}{l} wg_1g_2(w)^{-1} = wg_1(g_2(w))^{-1} = \\ (wg_2(w)^{-1})(g_2(w)g_1(g_2(w))^{-1}) \in N_m(W(\overline{b},\overline{y})). \end{array}$$

□

The braid group Br_m on m strings acts on $\mathbb{F}_m$. Below, we fix a set $\{a_1,\dots,a_{m-1}\}$ of so called *standard generators*, i.e., generators of Br_m being subject to the relations

$$\begin{array}{lll} a_ia_{i+1}a_i & = a_{i+1}a_ia_{i+1} & 1 \leq i \leq n-1, \\ a_ia_k & = a_ka_i & \mid i-k \mid \geq 2 \end{array}$$

and acting on $\mathbb{F}_m$ as follows:

$$\begin{array}{ll} a_i(x_j) = x_j & \text{if } i \neq j, j+1; \\ a_i(x_i) = x_ix_{i+1}x_i^{-1}; & \\ a_i(x_{i+1}) = x_i. & \end{array}$$

Therefore one can associate a set of relations $R_{W(B,\mathbb{F}_m)}$ to any subgroup $B \subset Br_m$. As in the case of the free groups, for $k \leq m$ the group Br_k is identified with the subgroup of Br_m generated by the first $k-1$ standard generators $a_1,\dots,a_{k-1}$, and we denote by $B_{k,i}$, $k+i \leq m$, the subgroup of Br_m generated by $a_{i+1},\dots,a_{i+k-1}$. Let $\Delta_{k,i}$ be the Garside element in the braid group $B_{k,i}$:

$$\Delta_{k,i} = (a_{i+1}\dots a_{i+k-1})\dots(a_{i+1}a_{i+2}a_{i+3})(a_{i+1}a_{i+2})a_{i+1}.$$

We have

$$\Delta_{k,i}(x_{i+j}) = (x_{i+1} \dots x_{i+k-j}) x_{i+k-j+1} (x_{i+1} \dots x_{i+k-j})^{-1} \qquad (7)$$

for $i+1 \leq j \leq i+k$ (we put here $x_{i+1} \dots x_{i+k-j} = \mathbf{1}$ if $j = i+k$) and

$$\Delta_k(x_j) = x_j$$

for $j \leq i$ and $j > i+k$. The Garside element in Br_k will be denoted by Δ_k.

Let $\mathrm{sh} = \mathrm{sh}_m : Br_m \to Br_{2m}$ be the homomorphism given by $\mathrm{sh}(a_j) = a_{m+j}$ for $j = 1, \dots, m-1$. For any subgroup B of Br_m, its image $\mathrm{sh}(B)$ will be called the *shift* of the group B by m.

Below, we will identify $\mathbb{F}_k$ with the fundamental group $\pi_1(D_k \setminus \{p_1, \dots, p_k\}, p_0)$, where $D_k = \{ z \in \mathbb{C} \mid |z| \leq k+1 \}$, $p_0 = -i(k+1)$, $p_j = -j$ for $j = 1, \dots, k$, and for each j the element x_j will be identified with the loop consisting of the segment $\gamma_j = \{ z \in \mathbb{C} \mid z = -i(k+1) - t(j + i(\varepsilon - k - 1)), 0 \leq t \leq 1 \}$ for some $0 < \varepsilon << 1$, the circuit around the circle $\{|z+j| = \varepsilon\}$ in the counterclockwise direction and the return along l_j (see Fig. 1). For so chosen base $x_1, \dots, x_k$, the standard generators a_j of Br_k, $j = 1, \dots, k-1$, are identified with half-twists defined by the segments $[-j-1, -j] = \{ z = -(j+1) + t \mid 0 \leq t \leq 1 \}$ (see details, for example, in [Moi-Te2]). The element $l_k = x_1 \dots x_k$ coincides in $\pi_1(D_k \setminus \{p_1, \dots, p_k\}, p_0)$ with circuit around the boundary ∂D_k of D_k

in the counterclockwise direction and it is fixed under the action of Br_k.



Fig. 1

We say that an ordered set $\{y_1, \dots, y_m\}$ consisting of elements of $\mathbb{F}_m$ is a *good geometric base* of $\mathbb{F}_m$ if there is $b \in Br_m$ such that $y_j = b(x_j)$ for $j = 1, \dots, m$. An ordered set $\{y_1, \dots, y_k\}$, $k \le m$, of elements of $\mathbb{F}_m$ is called *good* if the set $\{y_1, \dots, y_k\}$ can be extended to a good geometric base $\{y_1, \dots, y_m\}$ of $\mathbb{F}_m$. It is well-known that $\{y_1, \dots, y_m\}$ is a good geometric base of $\mathbb{F}_m$ if and only if each y_j is conjugated to some $x_{k(j)}$ in $\mathbb{F}_m$ and $y_1 \dots y_m = l_m$.

Consider an element $w(x_1, \dots, x_m) = x_{j_1}^{\varepsilon_1} \dots x_{j_n}^{\varepsilon_n}$, $\varepsilon_j = \pm 1$, in $\mathbb{F}_m$ of the letter length n. For $k > n$ and $p \ge 1$, the element

$$\mathrm{m}_{pmk,k,m}(w) = x^{\varepsilon_1}_{(pk-\varepsilon_1(k-1))m+j_1} \dots x^{\varepsilon_n}_{(pk-\varepsilon_n(k-n))m+j_n} \in \mathbb{F}_{m(p+1)k}$$

is called the *moving apart* of w by m with center at pmk.

Below, we will use the following claim which is obvious from geometric point of view.

Claim 1.7. *Let* $w_k(x_1, \dots, x_m) = x_{j_{k,1}}^{\varepsilon_{k,1}} \dots x_{j_{k,n_k}}^{\varepsilon_{k,n_k}}$, $\varepsilon_{k,j} = \pm 1$, $k = 1, 2$, *be two words in the letters* $x_1, \dots, x_m$ *and their inverses of the letter length*

n_k less than n. Then for each pair i_1, i_2 such that $1 \leq i_1, i_2 \leq m$, the pair $(y_1 = \overline{w}_1 x_{nm+i_1} \overline{w}_1^{-1}, y_2 = \overline{w}_2 x_{3nm+i_2} \overline{w}_2^{-1})$ is good in $\mathbb{F}_{4mn}$, where

$$\overline{w}_1 = \mathrm{m}_{mn,n,m}(w_1) = x^{\varepsilon_{1,1}}_{(n-\varepsilon_{1,1}(n-1))m+j_{1,1}} \dots x^{\varepsilon_{1,n_1}}_{(n-\varepsilon_{1,n_1}(n-n_1))m+j_{1,n_1}}$$

and

$$\overline{w}_2 = \mathrm{m}_{3mn,n,m}(w_1) = x^{\varepsilon_{2,1}}_{(3n-\varepsilon_{2,1}(n-1))m+j_{2,1}} \dots x^{\varepsilon_{2,n_2}}_{(3n-\varepsilon_{2,n_2}(n-n_2))m+j_{2,n_2}}.$$

Proof. The elements y_1 and y_2 are represented by loops of the form drawn in Fig. 2. Obviously, such pair (y_1, y_2) is good. $\square$



Fig. 2

Let $h : \{1, \dots, m\}^2 \to \mathbb{Z}$ be a non-negative integer-valued function. Put

$$I_h = \{(i, j, k) \in \mathbb{Z}^3 \mid 1 \leq i, j \leq m,\ 1 \leq k \leq h(i, j)\}$$

and let $f : I_h \to \mathbb{F}_m^2$ be a map to $\mathbb{F}_m^2 = \mathbb{F}_m \times \mathbb{F}_m$.

Claim 1.8. *For any map $f : I_h \to \mathbb{F}_m^2$, there are $M \in \mathbb{N}$, a finite set $I_{h,f} \subset \{1, \dots, M\}^2$ and a map $F : I_{h,f} \to \mathbb{F}_M^2$ such that*

- (i) *the pair $F(i, j) = (y_i, y_j)$ is a good pair in $\mathbb{F}_M$ for each $(i, j) \in I_{h,f}$;*
- (ii) *the sets of relations*

$$R_{W_f} = \{ x_i^{-1} w_{i,j,k} x_j w_{i,j,k}^{-1} = \mathbf{1} \mid (i, j, k) \in I_h,\ w_{i,j,k} = f(i, j, k) \}$$

and

$$R_{W_F} = \{ y_i^{-1} y_j = \mathbf{1} \mid (y_i, y_j) \in \operatorname{Im} F \}$$

are equivalent.

Proof. By Claim 1.7, there is an integer N such that the pairs

$$(x_i, \mathrm{m}_{Nm,N,m}(w_{i,j,k})x_{Nm+j}\mathrm{m}_{Nm,N,m}(w_{i,j,k})^{-1})$$

are good for all $(i,j,k) \in I_h$. Obviously, the set of relations R_{W_f} in $\mathbb{F}_m$ is equivalent to the union of two sets of relations $R_{\mathrm{m}(W_f)}$ and $R_{mod\, m}$ in $\mathbb{F}_{2Nm}$, where

$$R_{\mathrm{m}(W_f)} = \{x_i = \mathrm{m}_{Nm,N,m}(w_{i,j,k})x_{Nm+j}\mathrm{m}_{Nm,N,m}(w_{i,j,k})^{-1} \mid (i,j,k) \in I_h\}$$

and

$$R_{mod\, m} = \{x_i = x_j \mid i \equiv j \mod m)\}.$$

For each $i \leq m$ and each $k > 0$ the pair (x_i, x_{i+mk}) is good which completes the proof of Claim 1.8 if we put $M = 2Nm$ and $R_{W_F} = R_{\mathrm{m}(W_f)} \cup R_{mod\, m}$. □

By Claim 1.6, for a subgroup B of Br_m generated by $b_1, \dots, b_n$, the group $G(W(B, \mathbb{F}_m))$ is canonically isomorphic to

$$\begin{aligned} &G(W(\overline{b}, \overline{x})) = \\ &< x_1, \dots, x_m \mid x_i^{-1}b_j(x_i) = \mathbf{1},\ i = 1, \dots, m,\ j = 1, \dots, n > . \end{aligned} \tag{8}$$

Claim 1.9. *Presentation (8) defines on $G(W(\overline{b}, \overline{x}))$ a structure of a C-group.*

Proof. We have

$$b(x_i) = w_{b,i}(\bar{x})^{-1}x_j w_{b,i}(\bar{x}) \tag{9}$$

for some $j = \sigma_b(i)$ and $w_{b,i}(\bar{x}) \in \mathbb{F}_m$, where $\sigma_b \in \mathfrak{S}_m$ is the image of $b \in Br_m$ under the canonical epimorphism $\sigma : Br_m \to \mathfrak{S}_m$ with $\ker \sigma = P_m$ (we denote by $\mathfrak{S}_m$ the symmetric group acting on the set $\{1, \dots, m\}$ and by P_m the subgroup of pure braids). Therefore, the relations $x_i^{-1}b_j(x_i) = \mathbf{1}$ are equivalent to the C-relations $x_i = w_{b_j,i}(\bar{x})^{-1}x_j w_{b_j,i}(\bar{x})$. □

The following lemma is well known.

Lemma 1.10. *Let $b_k = g a_j^{k+1} g^{-1} \in Br_m$. Then the set of relations*

$$\{x_i^{-1}b_k(x_i) = \mathbf{1} \mid i = 1, \dots, m\}$$

in $\mathbb{F}_m$ is equivalent to one single relation:

(0) $g(x_j) = g(x_{j+1})$ *if* $k = 0$;
(1) *the commutant* $[g(x_j), g(x_{j+1})] = \mathbf{1}$ *if* $k = 1$ *or* -3;
(2) $g(x_j)g(x_{j+1})g(x_j) = g(x_{j+1})g(x_j)g(x_{j+1})$ *if* $k = 2$.

Proof. Consider a free base $\{y_1, \dots y_m\}$ of the group $\mathbb{F}_m$, where $y_i = g(x_i)$ for $i = 1, \dots, m$. By Claim 1.6, the sets of relations $\{x_i^{-1} b_k(x_i) = \mathbf{1} \mid i = 1, \dots, m\}$ and $\{y_i^{-1} b_k(y_i) = \mathbf{1} \mid i = 1, \dots, m\}$ are equivalent for each k.

We have

$$b_k(y_i) = g(x_i) = g a_j^{k+1} g^{-1}(g(x_i)) = g(x_i) = y_i$$

if $i \neq j, j+1$,

$$\begin{aligned} b_k(y_j) &= g a_j^{k+1}(x_j) = \\ &\begin{cases} g(x_j x_{j+1} x_j^{-1}) & \text{if } k = 0; \\ g((x_j x_{j+1}) x_j (x_j x_{j+1})^{-1}) & \text{if } k = 1; \\ (g((x_j x_{j+1} x_j) x_{j+1} (x_j x_{j+1} x_j)^{-1}) & \text{if } k = 2, \end{cases} \end{aligned}$$

and

$$\begin{aligned} b_k(y_{j+1}) &= g a_j^{k+1}(x_{j+1}) = \\ &\begin{cases} g(x_j) & \text{if } k = 0; \\ g(x_j x_{j+1} x_j^{-1}) & \text{if } k = 1; \\ g((x_j x_{j+1}) x_j (x_{j+1} x_j)^{-1}) & \text{if } k = 2. \end{cases} \end{aligned}$$

Therefore

$$b_k(y_j) = \begin{cases} y_j y_{j+1} y_j^{-1} & \text{if } k = 0; \\ (y_j y_{j+1}) y_j (y_j y_{j+1})^{-1} & \text{if } k = 1; \\ (y_j y_{j+1} y_j) y_{j+1} (y_j y_{j+1} y_j)^{-1} & \text{if } k = 2 \end{cases}$$

and

$$b_k(y_{j+1}) = \begin{cases} y_j & \text{if } k = 0; \\ y_j y_{j+1} y_j^{-1} & \text{if } k = 1; \\ (y_j y_{j+1}) y_j (y_{j+1} y_j)^{-1} & \text{if } k = 2. \end{cases}$$

Thus, for each k the set of relations $\{y_i^{-1} b_k(y_i) = \mathbf{1} \mid i = 1, \dots, m\}$ is reduced to one single relation

$$\begin{aligned} y_j y_{j+1}^{-1} &= \mathbf{1} \quad \text{if } k = 0; \\ y_j y_{j+1} y_j^{-1} y_{j+1}^{-1} &= \mathbf{1} \quad \text{if } k = 1; \\ (y_j y_{j+1} y_j)(y_{j+1} y_j y_{j+1})^{-1} &= \mathbf{1} \quad \text{if } k = 2. \end{aligned}$$

□

Claim 1.11. *Let y_1, y_2 be a good pair in $\mathbb{F}_m$. Then there is an element $b = g a_1 g^{-1} \in Br_m$ such that the set of relations $R_{W(<b>, \mathbb{F}_m)}$ is equivalent to the single relation*

$$R_b = \{y_1^{-1} y_2 = \mathbf{1}\}.$$

Proof. Since y_1, y_2 is a good pair, it can be included in a good geometric base $\{y_1, \dots, y_m\}$ of $\mathbb{F}_m$. Therefore there is an element $g \in Br_m$ such that $g(x_i) = y_i$ for $i = 1, \dots, m$. By Lemma 1.10, $R_{W(<b>,\mathbb{F}_m)}$ is equivalent to R_b. $\square$

Consider the braid group Br_{2m}. Put $a_{m,i} = a_i a_{2m-i}$,

$$\bar{c}_{m,i} = (a_{m,i}a_{m,i+1} \dots a_{m,m-1})a_m(a_{m,i}a_{m,i+1} \dots a_{m,m-1})^{-1} \tag{10}$$

for $1 \le i \le m-1$, $\bar{c}_{m,m} = a_m$, and

$$c_{m,i} = \Delta^{-1}_{m,m}\bar{c}_{m,i}\Delta_{m,m}. \tag{11}$$

Claim 1.12. *The set of relations $R_{W(<c_{m,i}>,\mathbb{F}_{2m})}$ is equivalent to the single relation*

$$R_{c_{m,i}} = \{\, x^{-1}_{m+i}x_i = \mathbf{1} \,\}.$$

Proof. Using (7), one can check that

$$\Delta^{-1}_{m,m}(a_{m,i}a_{m,i+1} \dots a_{m,m-1})(x_m) = x_i$$

and

$$\Delta^{-1}_{m,m}(a_{m,i}a_{m,i+1} \dots a_{m,m-1})(x_{m+1}) = x_{m+i}.$$

Therefore Claim 1.12 follows from (10), (11), and Claim 1.10. $\square$

For any subgroup B of Br_m, denote by $d(B) = d_m(B)$ (respectively, $\overline{d}(B) = \overline{d}_m(B)$) the subgroup of Br_{2m} generated by the elements of B, $\mathrm{sh}_m(B)$, and the elements $c_{m,1}, \dots, c_{m,m}$ (respectively, generated by the elements of B and the elements $c_{m,1}, \dots, c_{m,m}$), and call it the *doubling* (respectively, the *reduced doubling*) of the group B.

The doubling (respectively, the reduced doubling) of a group B can be iterated if we put

$$d^n(B) = d_{2^{n-1}m}(d^{n-1}(B)) \subset Br_{2^n m}. \tag{12}$$

Claim 1.13. *For any $n \in \mathbb{N}$ the following sets of relations*

$$R_{W(B,\mathbb{F}_m)},\ R_{d^n(W(B,\mathbb{F}_m))},\ R_{\overline{d}^n(W(B,\mathbb{F}_m))},\ R_{W(d^n(B),\mathbb{F}_{2^n m})},\ and R_{W(\overline{d}^n(B),\mathbb{F}_{2^n m})}$$

are equivalent.

Proof. It follows from Claims 1.4 and 1.12. $\square$

We say that a C-group G belongs to a subclass $\mathcal{C}_{Br}$ if it is C-isomorphic to a group given by the presentation

$$G =< x_1, \dots, x_m \mid R_{W(B,\mathbb{F}_m)} >$$

for some m and some finitely generated subgroup B of Br_m.

Theorem 1.14. *For any C-group G there are $M_G \in \mathbb{Z}$ and a subgroup B_G of Br_M generated by a finite set $\{b_1, \dots, b_{n_G}\}$ of elements conjugated in Br_M to the standard generator a_1 such that the group G is C-isomorphic to $G(W(B, \mathbb{F}_M))$.*

Proof. It follows from Claims 1.4, 1.7 1.8, 1.11, and 1.12. ☐

Corollary 1.15. *$\mathcal{C}_{Br} = \mathcal{C}$.*

Recall also the following theorem.

Theorem 1.16. ([Ar])

(i) *A C-group $G \in \mathcal{L}$ if and only if there are an integer m and an element $b \in Br_m$ such that G is C-isomorphic to*

$$G(W(<b>, \bar{x})) = < x_1, \dots, x_m \mid x_i = b(x_i) \text{ for } i = 1, \dots, m > . \tag{13}$$

(ii) *A C-group G with presentation* (13) *belongs to $\mathcal{K}$ if and only if the permutation $\sigma_b \in \mathcal{S}_m$ consists of one cycle of the length m, i.e., σ_b acts transitively on the set $\{1, \dots, m\}$.*

2. Full twist factorizations

It is well-known that the element (called the *full twist*)

$$\Delta_m^2 = (a_1 \dots a_{m-1})^m$$

is the generator of the center of Br_m. We have

$$\Delta_m^2(x_i) = (x_1 \dots x_m) x_i (x_1 \dots x_m)^{-1}. \tag{14}$$

In particular, the full twist Δ_m^2 leaves fixed the element $l_m = x_1 \dots x_m$.

Consider the braid group Br_{2m}. Put

$$\bar{r}_m = \bar{c}_{m,m} \bar{c}_{m,m-1} \dots \bar{c}_{m,1}$$

and

$$r_m = c_{m,m} c_{m,m-1} \dots c_{m,1},$$

where $\bar{c}_{m,i}$ and $c_{m,i}$ were defined by means of (10) and (11).

Lemma 2.1. *In the braid group Br_{2m}, we have*

$$\Delta_{2m} = \Delta_{m,0}^2 \Delta_{m,m}^2 \bar{r}_m. \tag{15}$$

Proof. Identify $Br_{2(m-1)}$ with subgroup $B_{2(m-1),1} \subset Br_{2m}$. By induction on m, we have

$$\bar{r}_m = \bar{c}_{m,m} \bar{c}_{m,m-1} \dots \bar{c}_{m,2} \bar{c}_{m,1} = \Delta_{(m-1),m}^{-2} \Delta_{m-1,1}^{-2} \Delta_{2(m-1),1} \bar{c}_{m,1}.$$

Therefore to prove Lemma 2.1, it is sufficient to show that the actions on $\mathbb{F}_{2m}$ of the elements Δ_{2m} and

$$b = \Delta_{m,0}^2 \Delta_{m,m}^2 \Delta_{m-1,m}^{-2} \Delta_{m-1,1}^{-2} \Delta_{2(m-1),1} \bar{c}_{m,1}$$

coincide.

We fix the base $\{x_1, \dots, x_{2m}\}$ of $\mathbb{F}_{2m}$ which is drawn in Fig. 1. Then the action of $\bar{c}_{m,1}$ on $\mathbb{F}_{2m}$ is induced by the half-twist defined by the path $\gamma_{m,1}$ which is drawn in Fig. 3.



Fig. 3

Therefore the element $y_1 = \bar{c}_{m,1}(x_1)$ is represented by the loop drawn in Fig. 4.



Fig. 4

The element $u_1 = \Delta_{2(m-1),1}(y_1)$ is represented by the loop drawn in Fig. 5.



Fig. 5

It is easy to see that $\Delta^{-2}_{m-1,m}\Delta^{-2}_{m-1,1}(u_1) = u_1$ and the element $v_1 = \Delta^{2}_{m,0}\Delta^{2}_{m,m}(u_1) = b(x_1)$ is represented by the loop drawn in Fig. 6.



Fig. 6

Therefore, by (7),

$$b(x_1) = \Delta_{2m}(x_1). \tag{16}$$

The element $y_i = \bar{c}_{m,1}(x_i)$, $i = 2, \dots, m$, is represented by the loop drawn in Fig. 7.



Fig. 7

The element $u_i = \Delta_{2(m-1),1}(y_i)$, $i = 2, \dots, m$, is represented by the loop drawn in Fig. 8.



Fig. 8

The element $v_i = \Delta^{-2}_{m-1,m}\Delta^{-2}_{m-1,1}(u_i)$, $i = 2, \dots, m$, is represented by the loop drawn in Fig. 9



Fig. 9

and the element $w_i = \Delta^2_{m,0}\Delta^2_{m,m}(v_i) = b(x_i)$, $i = 2, \dots, m$, is represented by the loop drawn in Fig. 10.



Fig. 10

Therefore

$$b(x_i) = \Delta_{2m}(x_i) \tag{17}$$

for $i = 2, \dots, m$.

It is easy to see that $\bar{c}_{m,1}(x_{2m-i}) = x_{2m-i}$ for $i = 1, \dots, m-1$. The element $y_{2m-i} = \Delta_{2(m-1),1}(x_{2m-i})$ is represented by the loop drawn in Fig. 11.



Fig. 11

The element $u_{2m-i} = \Delta^{-2}_{m-1,m}\Delta^{-2}_{m-1,1}(y_{2m-i})$ is represented by the loop drawn in Fig. 12



Fig. 12

and the element $v_{2m-i} = \Delta^2_{m,0}\Delta^2_{m,m}(u_{2m-i}) = b(x_{2m-i})$ is represented by the loop drawn in Fig. 13.



Fig. 13

Therefore, by (7),

$$b(x_{2m-i}) = \Delta_{2m}(x_{2m-i}) \tag{18}$$

for $i = 1, \ldots, m-1$.

The element $y_{2m} = \bar{c}_{m,1}(x_{2m})$ is represented by the loop drawn in Fig. 14.



Fig. 14

The element $u_{2m} = \Delta_{2(m-1),1}(y_{2m})$ is represented by the loop drawn in Fig. 15.



Fig. 15

It is easy to see that $\Delta^{-2}_{m-1,m}\Delta^{-2}_{m-1,1}(u_{2m}) = u_{2m}$ and the element $b(x_{2m}) = \Delta^2_{m,0}\Delta^2_{m,m}(u_{2m}) = x_1$. Therefore, by (7),

$$b(x_{2m}) = \Delta_{2m}(x_{2m}). \tag{19}$$

and Lemma 2.1 follows from (16) – (19). $\square$

Full twist formula for double number of strings.

$$\Delta^2_{2m} = \Delta^4_{m,0}\Delta^4_{m,m}r^2_m. \tag{20}$$

Proof. By Lemma 2.1,

$$\Delta_{2m} = \Delta^2_{m,0}\Delta^2_{m,m}\bar{r}_m.$$

It is easy to see that $[\Delta_{m,0}, \Delta_{m,m}] = \mathbf{1}$ and $[\Delta_{2m}, \Delta^2_{m,0}\Delta^2_{m,m}] = \mathbf{1}$. Therefore $[\bar{r}_{m,m}, \Delta^2_{m,0}\Delta^2_{m,m}] = \mathbf{1}$ and

$$\Delta^2_{2m} = \Delta^4_{m,0}\Delta^4_{m,m}\bar{r}^2_m.$$

Thus

$$\Delta^2_{2m} = \Delta^{-1}_{m,m}\Delta^2_{2m}\Delta_{m,m} = \Delta^4_{m,0}\Delta^4_{m,m}r^2_m.$$

$\square$

3. FACTORIZATION SEMIGROUPS OVER BRAID GROUPS

Let $\{g_i\}_{i\in I}$ be a set of elements of the braid group Br_m. For each $i \in I$ denote by $O_{g_i} \subset Br_m$ the set of the elements in Br_m conjugated to g_i (the orbit of g_i under the action of Br_m by the inner automorphisms). The union $U = \cup_{i\in I} O_{g_i} \subset Br_m$ is called the *full set of conjugates* of $\{g_i\}_{i\in I}$ and the pair (Br_m, U) an *equipped braid group*. In [Kh-Ku], a semigroup $S(Br_m, U)$, called a *factorization semigroup over* Br_m, was associated to each equipped braid group (Br_m, U). Recall that by definition, the semigroup $S(Br_m, U)$ is a semigroup generated by the alphabet U being subject to the relations

$$u_1 \cdot u_2 = u_2 \cdot (u_2^{-1} u_1 u_2) \quad \text{if } u_2 \neq \mathbf{1} \text{ and } u_1 \cdot \mathbf{1} = u_1 \quad \text{otherwise;}$$

$$u_1 \cdot u_2 = (u_1 u_2 u_1^{-1}) \cdot u_1 \quad \text{if } u_1 \neq \mathbf{1} \text{ and } \mathbf{1} \cdot u_2 = u_2 \quad \text{otherwise}$$

for all $u_1, u_2 \in U$.

There are two natural homomorphisms: the *product homomorphism* $\alpha = \alpha_U : S(Br_m, U) \to Br_m$ given by $\alpha(u) = u$ for each $u \in U$ and the homomorphism $\lambda : Br_m \to \mathrm{Aut}(S(Br_m, U))$ (the *conjugation action*) given by $\lambda(g)(u) = gug^{-1} \in U$ for all $g \in Br_m$. The action $\lambda(g)$ on $S(Br_m, U)$ is called the *simultaneous conjugation* by g. Put $\rho(g) = \lambda(g^{-1})$, $\lambda_S = \lambda \circ \alpha_U$ and $\rho_S = \rho \circ \alpha_U$. The orbit of an element $s \in S(Br_m, U)$ under the conjugation action of Br_m on $S(Br_m, U)$ is called the *type* of s.

Notice that $S : (Br_m, U) \mapsto (S(Br_m, U), \alpha_U, \lambda)$ is a functor from the category of equipped braid groups to the category of the semigroups over braid groups. In particular, if $U \subset V$ are two full sets of conjugates in Br_m, then the identity map $id : Br_m \to Br_m$ defines an embedding $id_{U,V} : S(Br_m, U) \to S(Br_m, V)$. So that, for each group Br_m, the semigroup $S_{Br_m} = S(Br_m, Br_m)$ is the *universal factorization semigroup over* Br_m, which means that each factorization semigroup $S_U = S(Br_m, U)$ over Br_m is canonically embedded in S_{Br_m} by id_{U,Br_m}.

Since $\alpha_U = \alpha_{Br_m} \circ id_{U,Br_m}$, there is no difference between the product homomorphisms α_U and α_{Br_m}, so the both are denoted simply by α.

For any s_1, $s_2 \in S(Br_m, U)$ we have

$$s_1 \cdot s_2 = s_2 \cdot \rho_S(s_2)(s_1) = \lambda_S(s_1)(s_2) \cdot s_1. \tag{21}$$

We say that an element $b \in Br_m$ has *the interlacing number* $l(b) = k$ if k is the smallest number such that b is conjugated in B_m to an element $\bar{b} \in B_{k,0}$, $b = g\bar{b}g^{-1}$. The element $\bar{b}$ will be called a *standard form* of b.

For each $s \in S_U$ denote by B_s the subgroup of Br_m generated by the images $\alpha(u_1), \dots, \alpha(u_n)$ of the elements $u_1, \dots, u_n$ of a factorization

$s = u_1 \cdot ... \cdot u_n$. It is easy to see that the subgroup B_s of Br_m does not depend on the presentation of $s \in S_U$ as a word in letters $u_i \in U$.

Associate to $s \in S_{B_m}$ one more group

$$\begin{array}{l} G(s) = G(W_s) = G(W(B_s, \mathbb{F}_m)) \simeq \\ < x_1, \dots, x_m \mid x_i^{-1} b(x_i) = \mathbf{1},\ i = 1, \dots, m,\ b \in B_s > . \end{array} \qquad (22)$$

Claim 3.1. *For $s_1, s_2 \in S_{Br_m}$*

(i) *there is a natural C-epimorphism*

$$\psi_{s_1} : G(s_2) \to G(s_1 \cdot s_2) \simeq G(s_2)/N_{G(s_2)}(\varphi_{W_{s_2}}(W_{s_1});$$

(ii) *if $s_2 = \lambda(b)(s_1)$, then there is a C-isomorphism $\gamma_b : G(s_1) \to G(s_2)$, in particular, if $s_1 = s_2$ then $G(s_1) = G(s_2)$ and $\gamma_{\mathbf{1}} = Id$.*

Proof. Straightforward. □

Let a factor u_i of $s = u_1 \cdot ... \cdot u_n$ have the interlacing number $l(u_i) = k_i$ and a standard form $\bar{u}_i \in Br_{k,0}$, $u_i = g_i \bar{u}_i g_i^{-1}$. Denote by $G_{u_i,loc}$ a subgroup of $G(s)$ generated by $\varphi_{W_s}(g_i(x_j))$, $j = 1, \dots, k_i$. It follows from (21) that for each factor u_i of $s = u_1 \cdot ... \cdot u_n$ the subgroup $G_{u_i,loc}$ is defined uniquely up to conjugation in $G(s)$.

The embedding $Br_m = B_{m,0} \subset Br_{2m}$ induces an embedding $S_{Br_m} \subset S_{Br_{2m}}$. Consider the homomorphism $\mathrm{sh} = \mathrm{sh}_m : Br_m \to Br_{2m}$ given by $\mathrm{sh}(a_i) = a_{m+i}$ for $i = 1, \dots, m-1$. It induces an embedding $\mathrm{sh} : S_{Br_m} \to S_{Br_{2m}}$. Put also

$$\widetilde{r}_m = c_{m,m} \cdot ... \cdot c_{m,1} \in S_{Br_{2m}},$$

where $c_{m,i}$ were defined in (11). Applying formula (20), we obtain

Full twist factorization formula for double number of strings. *For four elements $s_1, \dots, s_4 \in S_{Br_m}$ such that $\alpha(s_i) = \Delta_m^2$, the element*

$$\overline{s} = d(s_1, s_2, s_3, s_4) = s_1 \cdot s_2 \cdot sh(s_3) \cdot sh(s_4) \cdot \widetilde{r}_m \cdot \widetilde{r}_m \qquad (23)$$

is a factorization of Δ_{2m}^2 in $S_{Br_{2m}}$, i.e., $\alpha(\overline{s}) = \Delta_{2m}^2$.

If $s_1 = s_2 = s_3 = s_4 = s$ the doubling $d(s, s, s, s)$ will be denoted simply by

$$d(s) = d(s, s, s, s) = s \cdot s \cdot \mathrm{sh}(s) \cdot \mathrm{sh}(s) \cdot \widetilde{r}_m \cdot \widetilde{r}_m. \qquad (24)$$

and we put

$$d^{n+1}(s) = d(d^n(s)). \qquad (25)$$

Claim 3.2. *For $s \in S_{Br_m}$ such that $\alpha(s) = \Delta_m^2$, the groups $G(s)$ and $G(d(s))$ are C-isomorphic.*

Proof. It follows from Claim 1.13. □

Lemma 3.3. *Let* $s \in S_{Br_M}$ *be such that* $\alpha(s) = \Delta^2_M$. *Consider an element* $g \in Br_M$ *such that* $g(x_2) = x_i$ *for some* $i \leq M$. *Put* $y = g(x_1) \in \mathbb{F}_M$ *and* $b = g a_1 g^{-1}$. *Then*

(i) *the* C*-group* $\overline{G} = G(s)/N_{G(s)}(\varphi_{W_s}(x_i^{-1}y))$ *is canonically* C*-isomorphic to* $G(\overline{s})$*, where*
$$\overline{s} = d(\lambda_S(b)(d(s)), d(s), d(s), d(s)).$$

(ii) *the* C*-group* $\widetilde{G} = G(s)/N_{G(s)}(\varphi_{W_s}([x_i y]))$ *is canonically* C*-isomorphic to* $G(\widetilde{s})$*, where*
$$\widetilde{s} = d(\lambda_S(b^2)(d(s)), d(s), d(s), d(s)).$$

If $M = mk$ *and the set of relations* R_{W_s} *implies the relations* $R_{mod\, m}$ *in* $\mathbb{F}_M$*, then the set of relations* $R_{W_{\overline{s}}}$ *(respectively,* $R_{W_{\widetilde{s}}}$*) implies the set of relations* $R_{mod\, m}$ *in* $\mathbb{F}_{4M}$.

Proof. By Claim 3.2, the C-groups $G(s)$ and $G(d^2(s))$ are canonically C-isomorphic. By Claim 1.10, the set of relations $R_{W(<b^k>,\mathbb{F}_M)}$ is equivalent to the single relation $\{x_i^{-1}y = \mathbf{1}\}$ if $k = 1$ and $\{[x_i, y] = \mathbf{1}\}$ if $k = 2$. By Claim 3.1, the C-group $\overline{G}$ is canonically isomorphic to $G(b \cdot d^2(s))$ and the C-group $\widetilde{G}$ is canonically isomorphic to $G(b^2 \cdot d^2(s))$.

Put
$$s' = d(s) \cdot \mathrm{sh}_{2M}(d(s)) \cdot \mathrm{sh}_{2M}(d(s)) \cdot \widetilde{r}_{2M} \cdot \widetilde{r}_{2M}.$$
The element
$$\begin{aligned} & b^k \cdot d^2(s) = b^k \cdot d(s) \cdot s' = \lambda_S(b^k)(d(s)) \cdot b^k \cdot s' = \\ & \lambda_S(\lambda_S(b^k)(d(s)))(b^k) \cdot \lambda_S(b^k)(d(s)) \cdot s' = \\ & b^k \cdot \lambda_S(b^k)(d(s)) \cdot s' = \begin{cases} b \cdot \overline{s} \text{ if } k = 1, \\ b^2 \cdot \widetilde{s} \text{ if } k = 2, \end{cases} \end{aligned}$$
since
$$\alpha(d(s)) = \alpha(\lambda_S(b^k)(d(s))) = \Delta^2_{2M}$$
and the elements b^k and Δ^2_{2M} commute in Br_{2M}. Therefore, by Claim 3.1, there are the canonical epimorphisms $\psi_b : G(\overline{s}) \to \overline{G}$ and $\psi_{b^2} : G(\widetilde{s}) \to \widetilde{G}$. On the other hand, the factors $d(s) \cdot \mathrm{sh}_{2M}(d(s)) \cdot \mathrm{sh}_{2M}(d(s)) \cdot \widetilde{r}_{2M} \cdot \widetilde{r}_{2M}$ of the element $\overline{s}$ give the same set of relations as the element $d^2(s)$ gives. In particular, these relations imply the relations $\{x_{M+i} = x_i\} \in R_{mod\, M}$ in $\mathbb{F}_{4M}$ and, moreover, if the set of relations R_{W_s} implies the relations $R_{mod\, m}$ in $\mathbb{F}_M$, then the set of relations $R_{W_{\overline{s}}}$ (respectively, $R_{W_{\widetilde{s}}}$) implies the set of relations $R_{mod\, m}$ in $\mathbb{F}_{4M}$. The factor $\lambda_S(b^k)(c_{M,i}) = b^k c_{M,i} b^{-k}$ of $\lambda_S(b^k)(d(s))$ gives the relation $x_{M+i} = b c_{M,i} b^{-1}(x_{M+i})$ in $G(\overline{s})$ if $k = 1$ and $x_{M+i} = b^2 c_{M,i} b^{-2}(x_{M+i})$ in $G(\widetilde{s})$ if $k = 2$. But
$$b^k c_{M,i} b^{-k}(x_{M+i}) = b^k c_{M,i}(x_{M+i}) = b^k(x_i).$$

For $k=1,2$, we have $b(x_i)=y$ and $b^2(x_i)=yx_iy^{-1}$. The relations $\{x_{M+i}=y\}$ (respectively, $\{x_{M+i}=yx_iy^{-1}\}$) and $\{x_{M+i}=x_i\}$ imply the relation $x_i=y$ in $G(\overline{s})$ (respectively, $\{[x_i,y]=\mathbf{1}\}$ in $G(\widetilde{s})$). Therefore the canonical epimorphisms $\psi_b : G(\overline{s})\to\overline{G}$ and $\psi_{b^2} : G(\widetilde{s})\to\widetilde{G}$ are C-isomorphisms. □

Consider subclasses $\mathcal{C}_{\Delta^2,m}$, $m\in\mathbb{N}$, of the class $\mathcal{C}$ consisting of C-groups which are C-isomorphic to C-groups possessing C-presentations of the form

$$G=< x_1,\dots,x_m \mid x_i^{-1}w_{i,j,k}(\bar{x})^{-1}x_jw_{i,j,k}(\bar{x})=\mathbf{1},\ \ w_{i,j,k}(\bar{x})\in W>$$

for some finite set $W\subset\mathbb{F}_m$ such that the words $[x_i^{-1},x_1\dots x_m]$ belong to W for all $i=1,\dots,m$. Put

$$\mathcal{C}_{\Delta^2}=\bigcup_{m\in\mathbb{N}}\mathcal{C}_{\Delta^2,m}.$$

Claim 3.4. *Let $s\in S_{Br_m}$ be such that $\alpha(s)=\Delta_m^2$, then the C-group $G(s)\in\mathcal{C}_{\Delta^2,m}$. In particular, for $s=\Delta_m^2$ the group $G(\Delta_m^2)$ possesses the C-presentation*

$$G(\Delta_m^2)=< x_1,\dots,x_m \mid [x_i^{-1},x_1\dots x_m]=\mathbf{1}\ \textit{for}\ i=1,\dots,m > .$$

Proof. We have $\Delta_m^2(x_i)=(x_1\dots x_m)x_i(x_1\dots x_m)^{-1}$. Therefore, the relations $[x_i^{-1},x_1\dots x_m]=\mathbf{1}$, $i=1,\dots,m$, belong to the set of relations in presentation (22), since $\alpha(s)=\Delta_m^2$. □

Denote by $D_{k,m}$, $k\le m$, the full set of conjugates of the elements a_1 and Δ_k^2 in Br_m. Put $\mathcal{D}_{k,m}=S(Br_m,D_{k,m})$.

Theorem 3.5. *Let a C-group $G\in\mathcal{C}_{\Delta^2,m}$. Then there is $M\in\mathbb{N}$ and an element $s\in\mathcal{D}_{m,M}$ such that*

(i) *$\alpha(s)=\Delta_M^2$;*
(ii) *G and $G(s)$ are C-isomorphic.*

Proof. Let

$$G=< x_1,\dots,x_m \mid x_i^{-1}w_{i,j,k}(\bar{x})x_jw_{i,j,k}(\bar{x})^{-1}=\mathbf{1},\ \ w_{i,j,k}(\bar{x})\in W>,$$

for some finite set $W\subset\mathbb{F}_m$ such that the words $[x_i^{-1},x_1\dots x_m]$ belong to W for all $i=1,\dots,m$.

Numerate the words $w\in W$ so that the word $[x_i^{-1},x_1\dots x_m]$ has the number i. Denote by $G_n=\mathbb{F}_m/N_n$, where N_n is the normal closure of the set of words $\overline{W}_n=\{w_1,\dots,w_n\}$ in $\mathbb{F}_m$. For each $n\ge m$ we construct an element $s_n\in\mathcal{D}_{m,M_n}$ such that

(1) $\alpha(s_n)=\Delta_{M_n}^2$;
(2) G_n and $G(s_n)$ are C-isomorphic.

We start with $M_m = m$ and $s_m = \Delta^2_m \in \mathcal{D}_{m,m}$. By Claim 3.4, G_m and $G(s_m)$ are C-isomorphic.

Assume that for some $k \geq 0$ and $p \geq 0$ we have constructed an element $s_{m+k} \in \mathcal{D}_{m,M_{m+k}}$, where $M_{m+k} = 2^p m$, such that

(1) $\alpha(s_{m+k}) = \Delta^2_{M_{m+k}}$,
(2) G_{m+k} and $G(s_{m+k})$ are C-isomorphic,
(3) the set of relations $R_{W_{s_{m+k}}}$ implies the relations $R_{mod\, m}$.

Construct an element s_{m+k+1} having the similar properties as s_{m+k} has. Put $n = m + k$ and consider the word

$$w_{n+1} = x_{i_n}^{-1} w_{i_n,j_n,k_n}(\bar{x}) x_{j_n} w_{i_n,j_n,k_n}(\bar{x})^{-1}.$$

Let l be the letter length of the word w_{i_n,j_n,k_n}. Consider the element $\widetilde{s}_n = d^{l+2}(s_n)$. By Claim 1.13, the sets of relations $R_{W_{s_n}}$ and $R_{W_{d^{l+2}(s_n)}}$ are equivalent. Note that $\alpha(d^{l+2}(s_n)) = \Delta^2_{2^{l+2}M_n}$ and condition (3) is also realized for the element $\widetilde{s}_n = d^{l+2}(s_n)$. Put

$$y_n = \mathrm{m}_{2^l M_n,m}(w_{i_n,j_n,k_n}) x_{2^l M_n + j_n} \mathrm{m}_{2^l M_n,m}(w_{i_n,j_n,k_n})^{-1}.$$

By Claims 1.7 and 1.11, there is an element $b \in Br_{2^{l+2}M_n}$ conjugated to the standard generator a_1 and such that the set of relations $R(W_{<b>,\mathbb{F}_{2^{l+2}M_n}})$ is equivalent to the single relation $x_{i_n+2^{l+1}M_n} = y_n$. Applying Lemma 3.3 (i), the element

$$s_{n+1} = d(\lambda_S(b)(d(\widetilde{s}_n)), d(\widetilde{s}_n), d(\widetilde{s}_n), d(\widetilde{s}_n))$$

is a desired one which completes the proof of the Theorem. □

4. WEAK μ-EQUIVALENCE

Denote by $A_i = A_i(m)$ the full set of conjugates of the element a_1^{i+1} in Br_m, where $\{a_1, \dots, a_{m-1}\}$ is a set of standard generators of Br_m (recall that all generators $a_1, \dots, a_{m-1}$ are conjugated to each other in Br_m), and put $\mathcal{A}_k = S(Br_m, A_{\leq k})$, $\mathcal{A}^0_k = S(Br_m, A^0_{\leq k})$, where $A_{\leq k} = A_{-3} \cup (\cup_{i=-1}^k A_i)$ and $A^0_{\leq k} = \cup_{i=0}^k A_i$. We have the embedding $\mathcal{A}^0_k \subset \mathcal{A}_k$.

Below, we restrict our consideration to the case $k = 2$. The semigroup $\mathcal{A}^0_2$ is called the *braid monodromy cuspidal factorization semigroup* and $\mathcal{A}_2$ the *braid monodromy cuspidal factorization semigroup with negative nodes.*

Let $U \subset Br_m$ be a full set of conjugates containing $A_2 \cup A_{-3}$. Note that if $g \in A_1$ then $g^{-1} \in A_{-3}$. Consider a semigroup

$$\overline{S}_U = \langle\, u \in U) \;\mid\; R \in \mathcal{R} \cup \overline{\mathcal{R}}\,\rangle,$$

where $\mathcal{R}$ is the set of relations defining S_U (see section 3) and

$$\overline{\mathcal{R}} = \{\, g\cdot(g^{-1}) = (g^{-1})\cdot g = \mathbf{1} \mid g \in A_1 \,\}$$

is the set of *cancellation relations*. There is the canonical homomorphism of semigroups

$$c : \mathcal{S}_U \to \overline{S}_U .$$

We say that two elements s_1, $s_2 \in S_U$ are *weakly equivalent* if $c(s_1) = c(s_2)$. It is easy to see that $\overline{S}_U$ can be considered as a semigroup over Br_m with the product homomorphism $\overline{\alpha} : \overline{S}_U \to Br_m$ such that $\alpha = \overline{\alpha}\circ c$.

Let (s, μ_s) be a pair consisting of $s \in S_U$ and a homomorphism $\mu_s : G(s) \to \mathfrak{S}_N$ to the symmetric group $\mathfrak{S}_N$ for some N. We say that two pairs (s_1, μ_{s_1}) and (s_2, μ_{s_2}) are *equivalent* if s_1 and s_2 belong to the same orbit under the conjugation action of Br_m on S_U (i.e., there is $b \in Br_m$ such that $s_2 = \lambda(b)(s_1)$), and $\mu_2 = \mu_1 \circ \gamma_b$, where γ_b is defined in Claim 3.1.

Two pairs (s_1, μ_{s_1}) and (s_2, μ_{s_2}) is said to be obtained by *admissible transformation* from each other (notation: $(s_1, \mu_{s_1}) \leftrightarrow (s_2, \mu_{s_2})$) if there is $g \in A_1$ such that $s_2 = g\cdot(g^{-1})\cdot s_1$, $\mu_1 = \psi_{g\cdot(g^{-1})}\circ\mu_2$, and the order of the group $\mu_{s_2}(G_{g,loc})$ is greater than two. We say that two pairs $(s', \mu_{s'})$ and $(s'', \mu_{s''})$ are *weakly μ-equivalent* if there is a sequence of pairs $(s_1, \mu_1), \dots, (s_n, \mu_n)$ such that $(s', \mu_{s'}) = (s_1, \mu_1)$, $(s'', \mu_{s''}) = (s_n, \mu_n)$, and for each $i = 1, \dots, n-1$ the pairs (s_i, μ_i) and (s_{i+1}, μ_{i+1}) are either equivalent or can be obtained from each other by an admissible transformation.

Below, we consider the case then $U = \mathcal{A}_2$. Let $s = u_1 \cdot \ldots \cdot u_n \in \mathcal{A}_2$, $u_i \in A_{\leq 2}$. We say that a homomorphism $\mu_s : G(s) \to \mathfrak{S}_N$ is *generic* if

(i) μ_s is an epimorphism,
(ii) $\mu_s(\varphi_s(x_i))$ is a transposition in $\mathfrak{S}_N$ for each C-generator $\varphi_s(x_i)$ of the group $G(s)$,
(iii) the order of the group $\mu_s(G_{u_i,loc})$ is greater than two for each factor $u_i \in A_{-3} \cup A_1 \cup A_2$.

Let (s, μ_s) be a pair consisting of $s \in \mathcal{A}_2$ and a homomorphism $\mu_s : G(s) \to \mathfrak{S}_N$ for some N. By Claim 3.2, there is the canonical C-isomorphism $\psi_d : G(d(s)) \to G(s)$. It defines the pair $(d(s), d(\mu_s))$, where $d(\mu_s) = \mu_s \circ \psi_d$.

Claim 4.1. *Let $s \in \mathcal{A}_2$ and $\mu_s : G(s) \to \mathfrak{S}_N$ be a generic epimorphism. Then the homomorphism $d(\mu_s) : G(d(s)) \to \mathfrak{S}_N$ is generic.*

Proof. Straightforward. □

Theorem 4.2. *Let* $s = u_1 \cdot \ldots \cdot u_n \in \mathcal{A}_2 \subset S_{Br_m}$, $u_i \in A_{\leq 2}$, *such that* $\alpha(s) = \Delta_m^2$, *and* $\mu_s : G(s) \to \mathfrak{S}_N$ *be a generic epimorphism. Let* x_i *be a generator of* $\mathbb{F}_m$ *and* y *be an element conjugated in* $\mathbb{F}_m$ *to a generator* x_j *for some* j *such that* $\mu_s(\varphi_s(x_i))$ *and* $\mu_s(\varphi_s(y))$ *are two different commuting transpositions in* $\mathfrak{S}_N$. *Then there are* $M \in \mathbb{N}$ *and two pairs* $(\overline{s}, \mu_{\overline{s}})$ *and* $(\widetilde{s}, \mu_{\widetilde{s}})$, $\overline{s}$, $\widetilde{s} \in \mathcal{A}_2 \subset S_{Br_M}$, *such that*

- (i) $\alpha(\overline{s}) = \alpha(\widetilde{s}) = \Delta_M^2$,
- (ii) $G_{\overline{s}}$ *is* C*-isomorphic to* $G(s)$ *and* $G_{\widetilde{s}}$ *is* C*-isomorphic to the* C*-group* $G(s)/\{[\varphi_s(x_i), \varphi_s(y)] = \mathbf{1}\}$,
- (iii) *if* $[\varphi_s(x_i), \varphi_s(y)] \neq \mathbf{1}$, *then the types of* $\overline{s}$ *and* $\widetilde{s}$ *are different,*
- (iv) $\mu_{\overline{s}}$ *and* $\mu_{\widetilde{s}}$ *are generic epimorphisms onto* $\mathfrak{S}_N$,
- (v) *the pairs* $(\overline{s}, \mu_{\overline{s}})$ *and* $(\widetilde{s}, \mu_{\widetilde{s}})$ *are weakly* μ*-equivalent.*

Moreover, if $s \in \mathcal{A}_2^0 \subset S_{Br_m}$, *then* $\overline{s}$, $\widetilde{s} \in \mathcal{A}_2^0 \subset S_{Br_M}$.

Proof. By Claims 1.7, 1.8, 1.11, and 4.1, there are

1. $M_1 = 2^n m$ (for some $n \in \mathbb{N}$),
2. an element $y_1 \in \mathbb{F}_{M_1}$ which is a move apart of the element y such that (y_1, x_k) is a good pair for some $k \equiv i \mod m$,
3. an element $s_1 = d^n(s) \in \mathcal{A}_2 \subset S_{Br_{M_1}}$ such that $\mu_s(x_i) = \mu_{s_1}(x_k)$, $\mu_s(y) = \mu_{s_1}(y_1)$, where $\mu_{s_1} = d^n(\mu_s)$.

By definition of the good pairs, there is an element $g \in Br_{M_1}$ such that $g(x_1) = y_1$ and $g(x_k) = x_2$. Put $b = g a_1 g^{-1} \in Br_{M_1}$, $\overline{s} = d^2(s_1)$, and $\widetilde{s} = d(\lambda_S(b^2)(d(s)), d(s), d(s), d(s))$. By Claim 4.1, the groups $G(s)$ and $G(\overline{s})$ are canonically C-isomorphic. Therefore the generic epimorphism μ_s induces a generic epimorphism $\mu_{\overline{s}} : G(\overline{s}) \to \mathfrak{S}_N$. By Lemma 3.3 (ii) (see its proof), the groups $G(\widetilde{s})$ and $G(b^{-2} \cdot b^2 \cdot \overline{s})$ are canonically C-isomorphic to $G(\overline{s})/N_{G(\overline{s})}(\varphi_{W_{\overline{s}}}([x_j, y_1]))$, where $\widetilde{s} = d(\lambda_S(b^2)(d(s_1)), d(s_1), d(s_1), d(s_1))$. Therefore these isomorphisms induce generic epimorphisms $\mu_{b^{-2} \cdot b^2 \cdot \overline{s}} : G(b^{-2} \cdot b^2 \cdot \overline{s}) \to \mathfrak{S}_N$ and $\mu_{\widetilde{s}} : G(\widetilde{s}) \to \mathfrak{S}_N$. We have $(\overline{s}, \mu_{\overline{s}}) \leftrightarrow (b^{-2} \cdot b^2 \cdot \overline{s}, \mu_{b^{-2} \cdot b^2 \cdot \overline{s}})$, the pairs $(b^{-2} \cdot b^2 \cdot \overline{s}, \mu_{b^{-2} \cdot b^2 \cdot \overline{s}})$ and $(b^{-2} \cdot b^2 \cdot \widetilde{s}, \mu_{b^{-2} \cdot b^2 \cdot \widetilde{s}})$ are equivalent, and $(b^{-2} \cdot b^2 \cdot \widetilde{s}, \mu_{b^{-2} \cdot b^2 \cdot \widetilde{s}}) \leftrightarrow (\widetilde{s}, \mu_{\widetilde{s}})$. Therefore the pairs $(\overline{s}, \mu_{\overline{s}})$ and $(\widetilde{s}, \mu_{\widetilde{s}})$ are weakly μ-equivalent. This completes the proof of Theorem 4.2. □

5. Hurwitz curves

Let F_N be a relatively minimal ruled rational surface, $N \geq 1$, $\mathrm{pr} : F_N \to \mathbb{CP}^1$ the ruling, R a fiber of pr, and E_N the exceptional section, $E_N^2 = -N$. Below, we will identify $\mathrm{pr} : F_1 \to \mathbb{CP}^1$ with a linear projection $\mathrm{pr} : \mathbb{CP}^2 \to \mathbb{CP}^1$ with center at a point $p \in \mathbb{CP}^2$ (p is the blow down E_1 to the point).

Definition 5.1. *The image $\bar{H} = f(S) \subset F_N$ of a smooth map $f : S \to F_N \setminus E_N$ of an oriented closed real surface S is called a Hurwitz curve (with respect to* pr*) of degree m if there is a finite subset $Z \subset \bar{H}$ such that:*

(i) *f is an embedding of the surface $S \setminus f^{-1}(Z)$ and for any $s \notin Z$, $\bar{H}$ and the fiber $R_{pr(s)}$ of* pr *meet at s transversely and with positive intersection number;*

(ii) *for each $s \in Z$ there is a neighborhood $U \subset F_N$ of s such that $\bar{H} \cap U$ is a complex analytic curve, and the complex orientation of $\bar{H} \cap U \setminus \{s\}$ coincides with the orientation transported from S by f;*

(iii) *the restriction of* pr *to $\bar{H}$ is a finite map of degree m.*

For any Hurwitz curve $\bar{H}$ there is one and only one minimal $Z \subset \bar{H}$ satisfying the conditions from Definition 3.1. We denote it by $Z(\bar{H})$.

A Hurwitz curve $\bar{H}$ is called *cuspidal* if for each $s \in Z(\bar{H})$ there is a neighborhood U of s and local analytic coordinates z, w in U such that

(iv) $\mathsf{pr}_{|U}$ is given by $(z, w) \mapsto z$;

(v) $\bar{H} \cap U$ is given by $w^2 = z^k, k \geqslant 1$.

It is called *ordinary cuspidal* if $k \leq 3$ in (v) for all $s \in Z(\bar{H})$, and *nodal* if $k \leq 2$. Below, we will consider Hurwitz cuspidal curves $\bar{H}$ *with negative nodes* allowing $\bar{H}$ to have also singularities of the following form:

(vi) in a neighborhood $U \subset F_N$ of a point $s \in \bar{H}$ the intersection $U \cap \bar{H}$ splits into two branches meeting transversally at s with negative intersection number and meeting transversally the fibre $R_{\mathrm{pr}(s)}$ with positive intersection number.

Since $E_N \cap \bar{H} = \emptyset$, one can define a braid monodromy factorization $bmf(\bar{H}) \in S_{Br_m}$ of $\bar{H}$ as in the algebraic case (see, for example, [Kh-Ku]). To do this, we fix a fiber R_∞ meeting transversely $\bar{H}$ and consider the affine Hurwitz curve $H = \bar{H} \cap \mathbb{C}^2$, where $\mathbb{C}^2 = F_N \setminus (R_\infty \cup E_N)$. Choose $r >> 1$ such that $\mathrm{pr}(Z)$ belongs to the disc $D(r) = \{ \mid z \mid \leq r \} \subset \mathbb{C} = \mathbb{CP}^1 \setminus \mathrm{pr}(F_\infty)$, $Z = Z(\bar{H})$. Denote by $z_1, \dots, z_n$ the elements of $\mathrm{pr}(Z)$. Pick ε, $0 < \varepsilon << 1$, such that the discs $D_i(\varepsilon) = \{z \in \mathbb{C} \mid \mid z - z_i \mid \leq \varepsilon \}$, $i = 1, \dots, n$, are disjoint. Select arbitrary points $u_i \in \partial D_i(\varepsilon)$ and a point $u_0 \in \partial D(r)$. Choose disjoint simple paths $l_i \subset D(r) \setminus \bigcup_1^n D_i(r)$, $i = 1, \dots, n$, starting at u_0 and ending at u_i and renumber the points in a way that the product $\gamma_1 \dots \gamma_n$ of the loops $\gamma_i = l_i \circ \partial D_i(\varepsilon) \circ l_i^{-1}$ is equal to $\partial D(r)$ in $\pi_1(D(r) \setminus \{z_1, \dots, z_n\}, u_0)$.

Each γ_i, lifted to H, defines an element $b_i \in Br_m$. The factorization $b_1 \cdot \ldots \cdot b_n \in S_{Br_m}$ is called a *braid monodromy factorization* of $\bar{H}$. In fact, each b_i is conjugated to a braid monodromy standard form of some algebraic germ over z_i or $b_i \in A_{-3}$ if the corresponding singularity of $\bar{H}$ is a negative node. Hence, $bmf(\bar{H}) = b_1 \cdot \ldots \cdot b_n$ belongs to $\overline{\mathcal{P}} = S(Br_m, P \cup A_{-3})$, where P is the set of all braid monodromies of the germs of plane algebraic curves of degree m. The orbit of $bmf(\bar{H})$ under the conjugation action of Br_m on $\overline{\mathcal{P}}$ is called the *braid monodromy factorization type* of $\bar{H}$ and denoted by $bmt(\bar{H})$.

If $\bar{H}$ is an ordinary cuspidal Hurwitz curve with negative nodes, then its braid monodromy factorization $bmf(\bar{H}) \in \mathcal{A}_2$.

The following lemma is well known (see, for example, [Kh-Ku]).

Lemma 5.2. *For a Hurwitz curve $\bar{H} \subset F_N$ of degree m, we have*

$$\alpha(bmf(\bar{H})) = \Delta_m^{2N}.$$

The converse statement can be also proved in a straightforward way.

Theorem 5.3. ([Moi2]) *For any $b = b_1 \cdot \ldots \cdot b_n \in \overline{\mathcal{P}}$ such that $\alpha(b) = \Delta_m^{2N}$ there is a Hurwitz curve $\bar{H} \subset F_N$ with a braid monodromy factorization $bmf(\bar{H})$ equal to b.*

Definition 5.4. *Two Hurwitz curves $\bar{H}_1$ and $\bar{H}_2 \subset F_N$ are called H-isotopic if there is a fiberwise continuous isotopy $\phi_t : F_N \to F_N$, $t \in [0,1]$, smooth outside the fibers $R_{\mathsf{pr}(s)}$, $s \in Z(\bar{H}_1)$, and such that*

- (i) $\phi_0 = Id$;
- (ii) $\phi_t(\bar{H}_1)$ *is a Hurwitz curve for all* $t \in [0,1]$;
- (iii) $\phi_1(\bar{H}_1) = \bar{H}_2$;
- (iv) $\phi_t = Id$ *in a neighborhood of* E_N *for all* $t \in [0,1]$.

By Theorem 3.2 in [Kh-Ku], two cuspidal Hurwitz curves with negative nodes are H-isotopic if and only if they have the same braid monodromy factorization type.

Obviously, if $\bar{H}_1$ and $\bar{H}_2$ are H-isotopic, then the fundamental groups $\pi_1(F_N \setminus \bar{H}_1)$ and $\pi_1(F_N \setminus \bar{H}_2)$ are isomorphic.

Consider two sections $H_1 = \{u + 1 = 0,\, v = 0\}$ and $H_2 = \{u = 0,\, v = 0\}$ of the projection $\mathrm{pr}_1 : D \to D_1$ of a bi-disc $D = D_1 \times D_2$, where $D_1 = \{\sqrt{x^2 + y^2} \leq 2\}$ and $D_2 = \{\sqrt{u^2 + v^2} \leq 2\}$. Let h_t be an isotopy of H_1 given by equations

$$\begin{aligned} &u + 1 + 2t\rho(\sqrt{x^2 + y^2})(y + x^2 - 1) = 0, \\ &v - t\lambda\rho(\sqrt{x^2 + y^2})y = 0, \end{aligned}$$

where $\rho : \mathbb{R} \to \mathbb{R}$ is a monotone C^∞-function such that $\rho(s) = 0$ if $s \geq 2$ and $\rho(s) = 1$ if $s \leq 1$ and λ is a small positive constant. It is easy to check that

1. $h_t(H_1)$, $0 \leq t \leq 1$, and H_2 are symplectic curves in D with respect to the symplectic form $\omega = x \wedge y + u \wedge v$;
2. the sections $h_t(H_1)$ and H_2 have no common points if $0 \leq t < 1/2$;
3. the section $h_t(H_1)$ meets H_2 transversally at two points with, respectively, positive and negative intersection numbers if $1/2 < t \leq 1$.

We say that the "regular homotopy" $h_t(H_1) \cup H_2$ is the *creation of negative node*. The converse homotopy is called the *cancellation of negative node*.

Two Hurwitz curves $\bar{H}'$ and $\bar{H}'' \subset F_N$ are said to be obtained from each other by an *admissible transformation* if there is a neighborhood $U \subset F_N$ diffeomorphic to the bi-disc $D = D_1 \times D_2$ with local analytic coordinates $z = x + iy$ and $w = u + iv$ such that $\mathrm{pr}_{|U} = \mathrm{pr}_1 : D \to D_1$,

$$\bar{H}' \cap U = \{u + 1 = 0,\ v = 0\} \cup \{u = 0,\ v = 0\},$$

$$\bar{H}'' \cap U = \{u + 1 + 2(y + x^2 - 1)\rho = 0,\ v - \rho y = 0\} \cup \{u = 0,\ v = 0\},$$

and $\bar{H}' \cap (F_N \setminus U) = \bar{H}'' \cap (F_N \setminus U)$.

It is easy to see that if $\bar{H}'$ and $\bar{H}'' \subset F_N$ are obtained from each other by the above admissible transformation, then

$$bmf(\bar{H}'') = g \cdot (g^{-1}) \cdot bmf(\bar{H}') \tag{26}$$

for some $g \in A_1$.

We say that $\bar{H}'$ and $\bar{H}'' \subset F_N$ are *weakly equivalent* if there is a sequence of Hurwitz curves $\bar{H}_i$, $i = 1, \dots, n$, such that $\bar{H}' = \bar{H}_1$, $\bar{H}'' = \bar{H}_n$, and for each $i = 1, \dots, n-1$ the Hurwitz curves $\bar{H}_i$, $\bar{H}_{i+1}$ either are H-isotopic or are obtained from each other by an admissible transformation.

Claim 5.5. *Two cuspidal Hurwitz curves $\bar{H}'$ and $\bar{H}'' \subset F_N$ (possibly with negative nodes) are weakly equivalent if and only if their braid monodromy factorizations $bmf(\bar{H}')$ and $bmf(\bar{H}'')$ are weakly equivalent.*

Proof. It follows from (26) and Theorem 3.2 in [Kh-Ku]. □

Note that if two cuspidal Hurwitz curves $\bar{H}'$ and $\bar{H}'' \subset F_N$ are weakly equivalent then it is not necessary that their fundamental groups $\pi_1(F_N \setminus \bar{H}')$ and $\pi_1(F_N \setminus \bar{H}'')$ are isomorphic.

6. FUNDAMENTAL GROUPS OF THE COMPLEMENTS OF HURWITZ CURVES

The following theorem is well-known.

Zariski – van Kampen Theorem. *Let $\bar{H} \subset F_N$ be a Hurwitz curve of degree m having braid monodromy factorization $s = bmf(\bar{H})$. Then*

(i) *the fundamental group*

$$\pi_1(\mathbb{C}^2 \setminus \bar{H}) \simeq G(B_s),$$

where $\mathbb{C}^2 = F_N \setminus (E_N \cup R_\infty)$.

(ii) *the fundamental group*

$$\pi_1(\mathbb{CP}^2 \setminus \bar{H}) \simeq G(B_s)/\{\varphi_{W_s}(x_1 \dots x_m) = \mathbf{1}\}.$$

In the case $N = 1$, since $\alpha(s) = \Delta_m^2$ for $s = bmf(\bar{H})$, we get that the relations

$$[x_i^{-1}, x_1 \dots x_m] = \mathbf{1},\ i = 1, \dots, m \tag{27}$$

belong to the set of defining relations of presentation (22).

Denote by $\mathcal{H}$ the subclass of $\mathcal{C}$ consisting of the fundamental groups of the complements of the affine plane Hurwitz curves, i.e.,

$$\mathcal{H} = \{\, \pi_1(\mathbb{CP}^2 \setminus (\bar{H} \cup R_\infty))\, \},$$

where R_∞ is a generic line of the pencil defining the ruling $\mathrm{pr} : \mathbb{CP}^2 \to \mathbb{CP}^1$

Theorem 6.1. $\mathcal{H} = \mathcal{C}_{\Delta^2}$.

Proof. The inclusion $\mathcal{H} \subset \mathcal{C}_{\Delta^2}$ follows from (22), (27), and Claim 1.6.

The reverse inclusion follows from

Theorem 6.2. *For any C-group $G \in \mathcal{C}_{\Delta^2,m}$ there is a Hurwitz curve $\bar{H} \subset \mathbb{CP}^2$ with the singularities of the type $\{w^m = z^m\}$ such that $\pi_1(\mathbb{C}^2 \setminus H)$ is C-isomorphic to G, where $\mathbb{C}^2 = \mathbb{CP}^2 \setminus R_\infty$ and $H = \bar{H} \cap \mathbb{C}^2$.*

Proof. The braid monodromy of the singularity $\{w^m = z^m\}$ is equal to Δ_m^2. Therefore Theorem 6.2 follows from Theorems 3.5 and 5.3. □ □

Denote by $\mathcal{T}$ the subclass of $\mathcal{K}$ of the torus knot groups, that is, the fundamental groups $\pi_1(S^3 \setminus K_{p,q})$ of the knots

$$K_{p,q} = \{(z, w) \in S^3 \mid z = e^{2p\pi it},\, w = e^{2q\pi it},\, t \in [0, 1]\}$$

(p, q are co-prime, possibly $p = q = 1$), where

$$S^3 = \{\, (z, w) \in \mathbb{C}^2 \mid |z|^2 + |w|^2 = 2\, \}.$$

Theorem 6.3. $\mathcal{H} \cap \mathcal{K} = \mathcal{T}$.

Proof. Let $\bar{H} \subset \mathbb{CP}^2$ be a Hurwitz curve of degree m and G the fundamental group $\pi_1(\mathbb{C}^2 \setminus H)$ of the complement of its affine part $H \subset \mathbb{C}^2 = \mathbb{CP}^2 \setminus R_\infty$. It follows from (27) that the element $\varphi_{W_s}(x_1 \dots x_m)$ belongs to the center of the group G. Therefore, by Theorem 6.1 [Bu-Zi], $\mathcal{H} \cap \mathcal{K} \subset \mathcal{T}$.

The knot group $G_{p,q}$ of the knot $K_{p,q}$ has Wirtinger presentation

$$G_{p,q} =< x_1, \dots, x_p \mid x_i = (a_1 \dots a_{p-1})^q(x_i),\ i = 1, \dots, p > .$$

We have

$$(a_1 \dots a_{p-1})^{pq} = \Delta_p^{2q}.$$

Therefore there exists a Hurwitz curve $\widetilde{H} \subset F_q$ of degree p with braid monodromy factorization $bmf(\widetilde{H}) = s^p \in S_{Br_p}$, where $s = (a_1 \dots a_{p-1})^q$ is one of the generators of S_{Br_p} (the braid monodromy of the singularity given by the equation $w^p = z^q$). By Zariski – van Kampen Theorem, its fundamental group

$$\pi_1(\mathbb{C}^2 \setminus H) \simeq G_{p,q},$$

where $H = \mathbb{C}^2 \cap \widetilde{H}$ and $\mathbb{C}^2 = F_q \setminus R_\infty$, and R_∞ is a fibre of pr meeting $\widetilde{H}$ transversally.

As is known, see for example [Moi-Te],

$$\Delta_{p+1}^2 = \prod_{l=p+1}^{2} \prod_{k=1}^{l-1} b_{k,l}^2 = \Delta_p^2 \prod_{k=1}^{p} b_{k,p+1}^2,$$

where $b_{k,l} = (a_{l-1} \dots a_{k+1}) a_k (a_{l-1} \dots a_{k+1})^{-1}$ for $k < l$ (the notation $\prod_k^n$ states for from the left to the right product from k to n).

Put

$$s_1 = \prod_{k=1}^{p} (b_{k,p+1}^2) \in S_{Br_{p+1}},$$

where the product is taken in $S_{Br_{p+1}}$. The element $s^p \cdot s_1^q \in S_{Br_{p+1}}$ is a braid monodromy factorization of a Hurwitz curve in F_q, since $\alpha(s^p \cdot s_1^q) = \Delta_{p+1}^{2q}$. It is easy to see that this curve splits into two components one of which is H-isotopic to $\widetilde{H}$ and another one is a section $\widetilde{C}$ disjoint with E_q, $bmf(\widetilde{H} \cup \widetilde{C}) = s^p \cdot s_1^q$. The curves $\widetilde{H}$ and $\widetilde{C}$ meet transversally. Without loss of generality, we can assume that $\widetilde{C} \subset F_q$ is an algebraic section. It follows from the view of the braid monodromy factorization $bmf(\widetilde{H} \cup \widetilde{C}) = s^p \cdot s_1^q$ that the fundamental group $\pi_1(\mathbb{C}^2 \setminus (\widetilde{H} \cup \widetilde{C}))$ is canonically C-isomorphic to the direct product

$$\begin{array}{l} G_{p,q} \times \mathbb{F}_1 \simeq \\ < x_1, \dots, x_p \mid x_i = (a_1 \dots a_{p-1})^q(x_i),\ i = 1, \dots, p > \times < x_{p+1} > \end{array}$$

(the factor $\mathbb{F}_1 =< x_{p+1} >$ corresponds to the fundamental group $\pi_1(\mathbb{C}^2 \setminus \widetilde{C})$).

Choose non-homogeneous coordinates (u, v) in $\mathbb{C}^2 = F_q \setminus (E_q \cup R_\infty)$ such that $u = 0$ is an equation of $C = \widetilde{C} \cap (F_q \setminus (E_q \cup R_\infty))$, and coordinates (z, w) in the complement of some line $L = L_\infty \subset \mathbb{CP}^2$. Let $f : \mathbb{CP}^2 \to F_q$ be the cyclic (rational) covering of degree p given by $z = u^q$, $w = v$. It is branched along $\widetilde{C} \cup E_q$. Put $\bar{H} = f^{-1}(\widetilde{H})$ and $\bar{C} = F^{-1}(\widetilde{C})$. We have $L_\infty = f^{-1}(R_\infty \cup E_q)$, $\bar{H}$ is a Hurwitz curve of degree pq, and $\bar{C}$ is a line in $\mathbb{CP}^2$ meeting $\bar{H}$ transversally. It is easy to see that f_* gives the isomorphism of $\pi_1(\mathbb{C}^2 \setminus (\bar{H} \cup \bar{C}))$ and $G_{p,q} \times < x^q_{p+1} >$, where $< x^q_{p+1} >$ is generated by a circuit around $\bar{C}$. Therefore the canonical epimorphism

$$\psi : \pi_1(\mathbb{C}^2 \setminus (\bar{H} \cup \bar{C})) \to \pi_1(\mathbb{C}^2 \setminus \bar{H})$$

having $\ker \psi =< x^q_{p+1} >$ gives an isomorphism between $\pi_1(\mathbb{C}^2 \setminus \bar{H})$ and $G_{p,q}$. □

7. SYMPLECTOMORPHISMS OF GENERIC COVERINGS OF THE PLANE

An algebraic ramified covering of $\mathbb{CP}^2$ is a finite holomorphic map $f : X \to \mathbb{CP}^2$ of a normal projective irreducible complex surface X to the projective plane. The *ramification divisor* $R \subset X$ is the divisor of the jacobian of f (the multiplicity of R is the local degree of f minus 1). The *branch curve* $\bar{H} \subset \mathbb{CP}^2$ is the image of the support of R or, in other words, the set of points over which f is not locally invertible.

The fundamental group $\pi_1 = \pi_1(\mathbb{CP}^2 \setminus \bar{H}, p)$ of the complement of $\bar{H}$, where $p \in \mathbb{CP}^2 \setminus \bar{H}$, acts on the fibre $f^{-1}(p)$. Thus, a homomorphism (*monodromy of degree* $N = \deg f$) $\overline{\mu} = \overline{\mu}(f) : \pi_1 \to \mathfrak{S}_N$ from π_1 to the symmetric group $\mathfrak{S}_N$ is well-defined. The monodromy $\overline{\mu}$ is determined by f uniquely up to inner automorphism of the symmetric group. Conversely, by Grauert - Remmert theorem ([Gr-R]), a homomorphism $\overline{\mu} : \pi_1 \to \mathfrak{S}_N$ the image $Im\, \overline{\mu}$ of which acts transitively on a set consisting of N elements is the monodromy of some finite morphism $f : X \to \mathbb{CP}^2$ of $\deg f = N$.

To describe the fundamental group of the complement of a curve $\bar{H}$ of $\deg \bar{H} = m$, we fix a point $p \in \mathbb{CP}^2 \setminus \bar{H}$, choose a point $x \in \bar{H} \setminus Sing\, \bar{H}$, and consider a projective line $\Pi = \mathbb{CP}^1 \subset \mathbb{CP}^2$ meeting $\bar{H}$ transversally at x. Let $\Gamma \subset \Pi$ be a circle of small radius with center at x. The complex orientation on $\mathbb{CP}^2$ defines an orientation on Γ. Let γ be a loop consisting of a path L in $\mathbb{CP}^2 \setminus \bar{H}$, joining p with a point $q \in \Gamma$, the loop Γ (with positive direction) starting and ending at q, and a return to p along L in the opposite direction. Such loops γ (and the elements in π_1 corresponding them) are called *geometric generators* of the fundamental group $\pi_1 = \pi_1(\mathbb{CP}^2 \setminus \bar{H},\, p)$. It is well-known that π_1 is generated by the geometric generators, and if $\bar{H}$ is an irreducible curve then any two geometric generators are conjugated in π_1.

For each singular point s_i of multiplicity k_i of the curve $\bar{H}$, let us choose a small neighborhood $U_i \subset \mathbb{CP}^2$ such that $\bar{H} \cap U_i$ is given (in local analytic coordinates in U_i) by Weierstrass equation

$$w^{k_i} + \sum_{j=0}^{k_i-1} c_j(z) w^j = 0$$

(in particular, $w^2 = z^3$ if s_i is a cusp and $w^2 = z^2$ if s_i is a node). Let p_i be an arbitrary point in $U_i \setminus \bar{H}$. Recall that if s_i is a cusp then $\pi_1(U_i \setminus \bar{H}, p_i)$ is isomorphic to the braid group Br_3 on three strings and generated by two geometric generators (say, a_1 and a_2) satisfying the relation $a_1a_2a_1 = a_2a_1a_2$, and if s_i is a node then $\pi_1(U_i \setminus \bar{H}, p_i)$ is isomorphic to $\mathbb{Z} \oplus \mathbb{Z}$ and generated by two commuting geometric generators.

Choose smooth paths L_i in $\mathbb{CP}^2 \setminus \bar{H}$, connecting the points p_i with p. This choice defines homomorphisms $\psi_i : \pi_1(U_i \setminus \bar{H}, p_i) \to \pi_1$. We call $\psi_i(\pi_1(U_i \setminus \bar{H}, p_i)) = G_i$ *the local fundamental group* of the singular point s_i. The local fundamental groups G_i depend on the choice of the paths L_i and therefore are defined uniquely up to conjugation in π_1. If $bmf(\bar{H}) = s = u_1 \cdot \ldots \cdot u_n \in S_{Br_m}$ and the factor u_i corresponds to the singular point s_i of $\bar{H}$, then, in view of Zariski – van Kampen Theorem, the group G_i coincides (up to conjugation) with the image of $G_{u_i,loc}$ under the natural epimorphism $r : G(B_s) \to G(B_s)/\{\varphi_{W_s}(x_1 \ldots x_m) = \mathbf{1}\}$. The monodromy $\overline{\mu} : \pi_1 \to \mathfrak{S}_N$ of a generic covering $f : X \to \mathbb{CP}^2$ branched along $\bar{H}$ defined a homomorphism $\mu : G(s) \to \mathfrak{S}_N$, where $\mu = \overline{\mu} \circ r$ and $s = bmf(\bar{H})$, which we also call the *monodromy* of f.

A finite morphism $f : X \to \mathbb{CP}^2$, $\deg f = N$, ramified over a cuspidal curve $\bar{H}$ is called *a generic covering of the plane with discriminant curve* $\bar{H}$ if the monodromy μ of f is a generic epimorphism, i.e., it satisfies the following conditions:

(i) μ is an epimorphism,
(ii) the image $\mu(\gamma)$ of each geometric generator γ is a transposition in $\mathfrak{S}_N$,
(iii) the images $\mu(G_i)$ of the local fundamental groups G_i corresponding to the singular points of $\bar{H}$ are subgroups of $\mathfrak{S}_N$ of orders greater than 2.

The last condition is equivalent to the condition that X is a non-singular algebraic surface (see, for example, [Ku]). Note that if $\bar{H}$ is the discriminant curve of some generic covering, then its degree is an even number. For a generic covering f, the total transform $f^*(\bar{H})$ splits in the sum $f^*(\bar{H}) = 2R + C$, where $R \subset X$ is the ramification divisor of f having ramification multiplicity equal to two and C is a reduced curve in which f is not ramified. The restriction $f_{|R}$ to R is a morphism of degree 1.

In the symplectic case, by an observation of Gromov, the discussion of generic coverings of the plane given in terms of monodromies of generic coverings can be generalized to the symplectic situation as well (see Lemma 1 in [Go] and Proposition 10 in [Au] for more details). Firstly, given a cuspidal symplectic surface $\bar{H}$ (possibly with negative nodes) in $\mathbb{CP}^2$ and a generic epimorphism (monodromy) $\overline{\mu}$ of the fundamental group of its complement onto $\mathfrak{S}_N$, one can construct a smooth ramified covering $f : X \to \mathbb{CP}^2$ of degree N branched over $\bar{H}$. The preimage of the symplectic form vanishes on the normal bundle to the ramification surface $R \subset X$ and at the preimages of cusps in R. However, it is possible to add an exact 2-form $\eta \in \Lambda^2(X)$ so that $f^*\omega_{\mathbb{CP}^2} + \eta$ is symplectic. A subtle point is that, although there is no canonical way to lift the symplectic *form* to X, the resulting symplectic *structure* is defined canonically by a Moser-type argument.

We state the result in the form of a lemma.

Lemma 7.1. *Suppose that $\bar{H} \subset \mathbb{CP}^2$ is a cuspidal symplectic surface (possibly with negative nodes) and $\overline{\mu} : \pi_1(\mathbb{CP}^2 \setminus \bar{H}) \to \mathfrak{S}_n$ is a generic monodromy. Then there exists a unique symplectic four-manifold X with a smooth map $f : X \to \mathbb{CP}^2$ branched over $\bar{H}$ as described above such that the symplectic structure on X is the pull-back of the standard symplectic structure on $\mathbb{CP}^2$.*

Clearly, given a symplectic isotopy $\bar{H}_t$, $t \in [0, 1]$, of cuspidal symplectic surfaces such that there exists an epimorphism $\overline{\mu} : \pi_1(\mathbb{CP}^2 \setminus \bar{H}_0) \to \mathfrak{S}_N$, we obtain a family of symplectomorphic symplectic 4-manifolds X_t. (Recall that, by the definition of isotopy, the topology of $\mathbb{CP}^2 \setminus \bar{H}_t$ remains unchangeable.) Applying similar arguments as in [Kh-Ku], one can show that any cuspidal symplectic surface with negative nodes in $\mathbb{CP}^2$ is symplectically isotopic to a cuspidal symplectic Hurwitz curve with negative nodes. So in symplectic case to investigate generic covering of $\mathbb{CP}^2$ branched along a cuspidal symplectic surface $\bar{H}$, we can restrict ourselves to the case when $\bar{H}$ is a Hurwitz curve.

Consider two generic coverings $f' : X' \to \mathbb{CP}^2$ and $f'' : X'' \to \mathbb{CP}^2$ of degree N branched, respectively, along cuspidal Hurwitz curves (possibly with negative nodes) $\bar{H}'$ and $\bar{H}''$. Let $s' = bmf(\bar{H}')$ and $s'' = bmf(\bar{H}'')$ be braid monodromy factorizations of $\bar{H}'$ and $\bar{H}''$, and f' (respectively, f'') defined by the monodromy $\mu' : G(s') \to \mathfrak{S}_N$ (respectively, μ'') We say that $\bar{H}'$ and $\bar{H}''$ are *weakly μ-equivalent* if the pairs $(s', \mu_{s'})$ and $(s'', \mu_{s''})$ are weakly μ-equivalent.

Lemma 7.2. *Let two Hurwitz curves $\bar{H}'$ and $\bar{H}''$ be the branch curves of two generic coverings f' and f'' of the plane. If $\bar{H}'$ and $\bar{H}''$ are weakly μ-equivalent, then there is a symplectomorphism $h : X' \to X''$ such that the coverings f' and $f'' \circ h$ are regular homotopic.*

Proof. It follows from the definition of weak μ-equivalence Claim 5.5, and Theorem 3.3 in [Kh-Ku]. □

In algebraic case, if (X, L) is a polarized projective surface, that is, L is an ample line bundle on X, then, for sufficiently large k, the sections of $L^{\otimes k}$ define an embedding of X into some projective space $\mathbb{CP}^r$. The restriction of a generic projection $\mathbb{CP}^r \to \mathbb{CP}^2$ gives rise to a generic ramified covering $f_k : X \to \mathbb{CP}^2$ with branch curve $\bar{H}_k$. Observe that the degree of this covering equals $k^2 \deg L = k^2 c_1(L)^2$. The space of generic projections is path-connected and therefore we can give the following definition.

Definition 7.3. *The k-th braid monodromy invariant $\mu_k(X, L)$ of the polarized complex surface (X, L) is the type of the braid monodromy factorization of $\bar{H}_k$.*

The polarization canonically defines a symplectic structure ω_L on X. To see this, pick a positive $(1,1)$-form representing the first Chern class $c_1(L) \in H^2(X, \mathbb{Z})$. Every two such forms can be joined by a linear homotopy, hence they are diffeomorphic by Moser's theorem.

Theorem 7.4. *Let (X, L) and (X', L') be polarized projective surfaces. Suppose that $\mu_k(X, L) = \mu_k(X', L')$ for some k such that $k^2 c_1(L)^2 \geqslant 12$. Then (X, ω_L) and $(X', \omega_{L'})$ are symplectomorphic.*

Proof. By assumption, the branch curves $\bar{H}$ and $\bar{H}'$ of generic projections onto $\mathbb{CP}^2$, defined by sections of $L^{\otimes k}$ and $(L')^{\otimes k}$, respectively, have the same type of braid monodromy factorizations. Thus, by Theorem 3.3 in [Kh-Ku], $\bar{H}$ and $\bar{H}'$ are symplectically isotopic as cuspidal Hurwitz curves. By virtue of Theorem 3 in [Ku2], there exists a unique generic monodromy $\mu : \pi_1(\mathbb{CP}^2 \setminus \bar{H}) \to \mathfrak{S}_N$ for $N = k^2 c_1(L_1)^2$. Therefore, by Lemma 7.1, we have a unique family of symplectic ramified coverings of $\mathbb{CP}^2$ connecting X and X'. It follows from Moser's theorem that X and X' are symplectomorphic with the pulled back symplectic structures. However, these pull-backs are proportional to the symplectic structures defined by the polarizations with the same coefficient k, which completes the proof. □

Let (X,ω) be a compact symplectic 4-manifold with symplectic form ω whose class $[\omega]\in H^2(X,\mathbb{Z})$. Fix an ω-compatible almost complex structure J and corresponding Riemannian metric g. Let L be a line bundle on X whose first Chern class is $[\omega]$. By [Au], for $k >> 0$, the line bundle $L^{\otimes k}$ admits many approximately holomorphic sections so that one can choose three of them which give an approximately holomorphic generic covering $f_k : X\to\mathbb{CP}^2$ of degree $N_k = k^2\omega^2$ branched alone a cuspidal Hurwitz curve $\bar H$ (possibly with negative nodes). Denote, as above, by $bmf(\bar H_k)=s_k$ a braid monodromy factorization of the Hurwitz curve $\bar H_k$ and $\mu_k : G(B_{s_k})\to\mathfrak{S}_{N_k}$ the monodromy of f_k.

Definition 7.5. *The k-th braid monodromy invariant* $\overline{\mu}_k(X,\omega)$ *of a symplectic 4-manifold* (X,ω) *is the pair* $(bmf(\bar H_k),\mu_k)$.

Theorem 7.6. ([Au-Ka]) *If two symplectic 4-manifolds* (X',ω') *and* (X'',ω'') *are symplectomorphic, then, for k sufficiently large, the braid monodromy invariants* $\mu_k(X',\omega')$ *and* $\mu_k(X'',\omega'')$ *are weakly* μ*-equivalent.*

Weak equivalence appears in this statement, since the construction of an isotopy of the coverings depends continuously on a choice of tamed almost complex structures. Therefore, for symplectomorphic 4-manifolds, we obtain a "regular homotopy" of branch curves, which leads to the weak μ-equivalence of their braid monodromy factorizations. The following statement shows that in general case we can not avoid the appearance of negative nodes.

Theorem 7.7. *For any* $N\geq 4$*, there are two isotopic generic coverings* $f_1 : X\to\mathbb{CP}^2$ *and* $f_2 : X\to\mathbb{CP}^2$ *of degree N branched along cuspidal Hurwitz curves* $\bar H_1$ *and* $\bar H_2$ *without negative nodes such that*

(i) $\bar H_1$ *and* $\bar H_2$ *have different braid monodromy factorization types;*
(ii) *The pairs* $(bmf(\bar H_1),\mu_1)$ *and* $(bmf(\bar H_2),\mu_2)$ *are weakly* μ*-equivalent.*

Proof. Let $\bar H_N$ be the branch curve of a generic linear projection $\mathrm{pr} : X_N\to\mathbb{CP}^2$ of a smooth projective surface $X_N\subset\mathbb{CP}^3$ of $\deg X_N = N$. Denote by $m=N(N-1)$ the degree of $\bar H_N$ and $s=s_N=bmf(\bar H_N)$ its braid monodromy factorization. The projection pr defines a generic epimorphism (monodromy) $\mu=\mu_N : G(B_s)\to\mathfrak{S}_N$. By [Moi1], the C-group

$$\begin{aligned} &G(s)=G(W(B_s,\mathbb{F}_m))\simeq \\ &< x_1,\dots,x_m \mid x_i^{-1}b(x_i)=\mathbf{1},\ i=1,\dots,m,\ b\in B_s > \end{aligned} \tag{28}$$

is C-isomorphic to the braid group Br_N and μ coincides with canonical epimorphism $\sigma : Br_N \to \mathfrak{S}_N$ with the group of pure braids as its kernel. Without loss of generality, we can assume that $\varphi_s(x_1) = a_1 \in Br_N$. Since μ is an epimorphism, there is an element y conjugated in $\mathbb{F}_m$ to some generator x_i such that $\varphi_s(y) = a_2^{-2}a_3a_2^2$. Note that $[\varphi_s(x_1), \varphi_s(y)] \neq \mathbf{1}$ in Br_N, but $\mu(\varphi_s(x_1)) = (1,2)$ and $\mu(\varphi_s(y)) = (3,4)$ are two different commuting transpositions in $\mathfrak{S}_N$. Then, by Theorem 4.2, there are $M \in \mathbb{N}$ and two pairs $(\overline{s}, \mu_{\overline{s}})$ and $(\widetilde{s}, \mu_{\widetilde{s}})$, $\overline{s}$, $\widetilde{s} \in \mathcal{A}_2 \subset S_{Br_M}$, such that

1. $\alpha(\overline{s}) = \alpha(\widetilde{s}) = \Delta_M^2$,
2. $G_{\overline{s}}$ is C-isomorphic to $G(s) \simeq Br_N$ and $G_{\widetilde{s}}$ is C-isomorphic to the C-group $Br_N/\{[a_1, a_2^{-2}a_3a_2^2] = \mathbf{1}\}$,
3. the types of $\overline{s}$ and $\widetilde{s}$ are different,
4. $\mu_{\overline{s}}$ and $\mu_{\widetilde{s}}$ are generic epimorphisms onto $\mathfrak{S}_N$,
5. the pairs $(\overline{s}, \mu_{\overline{s}})$ and $(\widetilde{s}, \mu_{\widetilde{s}})$ are weakly μ-equivalent
6. $\overline{s}$, $\widetilde{s} \in \mathcal{A}_2^0 \subset S_{Br_M}$.

Now by Theorem 5.3, there are two cuspidal Hurwitz curves $\bar{H}_1$ and $\bar{H}_2$ in $\mathbb{CP}^2$ of degree M with $bmf(\bar{H}_1) = \overline{s}$ and $bmf(\bar{H}_2) = \widetilde{s}$. The generic epimorphisms $\mu_{\overline{s}}$ and $\mu_{\widetilde{s}}$ define two generic coverings $f' : X' \to \mathbb{CP}^2$ and $f'' : X'' \to \mathbb{CP}^2$ branched, respectively, along $\bar{H}_1$ and $\bar{H}_2$. By Lemma 7.2, there is a symplectomorphism $h : X' \to X''$ such that the coverings $f_1 = f' : X = X' \to \mathbb{CP}^2$ and $f_2 = f'' \circ h : X \to \mathbb{CP}^2$ are isotopic, but their branch Hurwitz curves have different braid monodromy factorization types. □

References

[Ar] Artin, E.: *Theorie der Zopfe.* Hamburg Abh. **4**, (1925), 47–72,

[Au] Auroux D.: *Symplectic 4-manifolds as branched coverings of $\mathbb{P}^2$.* Invent. Math. **139** (2000), 551–602.

[Au-Ka] Auroux D., Katzarkov L.: *Branched coverings of $\mathbb{P}^2$ and invariants of symplectic 4-manifolds.* Invent. Math. **142** (2000), 631–673.

[Bu-Zi] Burde G., Zieschang H.: *Knots.* Walter de Gruyter, Berlin · New York, (1985).

[Go] Gompf R.: *Symplectically aspherical manifolds with nontrivial π_2.* Math. Res. Letters **5** (1998), 599–603.

[Gr-R] Grauert H., Remmert R.: *Komplexe Raume.* Math. Ann., **136** (1958), 245-318.

[Kh-Ku] Kharlamov V., Kulikov Vik.S.: *On Braid Monodromy Factorizations.* Izv. Math. **67:3** (2003); see also `ArXiv:mathAG/0302113`

[Ku-Ku] Kulikov V.S., Kulikov Vik.S.: *Generic coverings of the plane with A-D-E- singularities.* Izv. Math. **64:6** (2000), 1153-1195.

[Ku-Te] Kulikov Vik. S., Teicher M.: *Braid monodromy factorization and diffeomorphism types.* Izv. Math. **64:2** (2000), 311-341.

[Ku] Kulikov Vik.S.: *On Chisini's Conjecture.* Izv. Math., 63:6, (1999), 1139-1170.

[Ku1] Kulikov Vik. S.: *Geometric realization of C-groups.* Izv. Math. **45**, No. 1 (1995), 197-206.

[Ku2] Kulikov Vik.S.: *Generalized Chisini's Conjecture.* Proc. of Steklov Math. Ins., vol. **241** (2003).

[Nem] Nemirovski S.: *On Kulikov's theorem on the Chisini Conjecture.* Izv. Math. **65:1**, (2001), 71-74.

[Moi1] Moishezon B.: *Stable branch curves and braid monodromies.* Springer L.N.M. **862** (1981), 107-192.

[Moi2] Moishezon B.: *The arithmetic of braids and a statement of Chisini.* Contemporary Math. **164** (1994), 151-175.

[Moi-Te] Moishezon B., Teicher M. *Braid group technique in complex geometry, I: line arrangements in $\mathbb{CP}^2$.* Contemporary Math. **78** (1988), 425-555.

[Moi-Te2] Moishezon B., Teicher M. *Braid group technique in complex geometry, V: The fundamental group of a complement of a branch curve of a Veronese generic projection.* Communications in Analysis and Geometry **4**, No. 11 (1996), 1-120.

Steklov Mathematical Institute, Gubkina str., 8, 119991 Moscow, Russia

E-mail address: `kulikov@mi.ras.ru`